

Lecture 5: Chapter 5

C C Moxley

UAB Mathematics

24 September 15

§5.1 Differences Between Statistics and Probability

In Chapters 2 and 3, we collected sample data and summarized the data to approximate certain probabilities/chances that certain outcomes would be observed. This is a statistical approach.

§5.1 Differences Between Statistics and Probability

In Chapters 2 and 3, we collected sample data and summarized the data to approximate certain probabilities/chances that certain outcomes would be observed. This is a statistical approach.

In Chapter 4, we determined expected chances of certain outcomes using a probabilistic approach.

§5.1 Differences Between Statistics and Probability

In Chapters 2 and 3, we collected sample data and summarized the data to approximate certain probabilities/chances that certain outcomes would be observed. This is a statistical approach.

In Chapter 4, we determined expected chances of certain outcomes using a probabilistic approach.

In this chapter, we combine the two methods! We use a statistical approach to estimate parameters used in a probabilistic approach.

§5.2 Probability Distributions

Definition (Random Variable)

A **random variable** is a variable (usually represented by x) that has a single numerical value, determined by chance, for each outcome of a procedure.

§5.2 Probability Distributions

Definition (Random Variable)

A **random variable** is a variable (usually represented by x) that has a single numerical value, determined by chance, for each outcome of a procedure.

Definition (Probability Distribution)

A **probability distribution** is a function that gives the probability for each value of the random variable. It is often described in a table, formula, or graph.

§5.2 Probability Distributions

Definition (Random Variable)

A **random variable** is a variable (usually represented by x) that has a single numerical value, determined by chance, for each outcome of a procedure.

Definition (Probability Distribution)

A **probability distribution** is a function that gives the probability for each value of the random variable. It is often described in a table, formula, or graph.

Definition (Continuous vs. Discrete Random Variables)

A **discrete random variable** can take on a value from a finite or countably infinite collection of values whereas a **continuous random variable** can take on a value from a collection of uncountably infinite values.

§5.2 Probability Distribution Requirements

A function P must satisfy the following conditions to be a probability distribution:

- P must map the random variable into probabilities, i.e. $P(x)$ should be a number between 0 and 1.

§5.2 Probability Distribution Requirements

A function P must satisfy the following conditions to be a probability distribution:

- P must map the random variable into probabilities, i.e. $P(x)$ should be a number between 0 and 1.
- $\sum P(x) = 1$, where we sum over all possible values of x . Sometimes rounding errors will cause these sums to be slightly above or below 1.

§5.2 Probability Distribution Requirements

A function P must satisfy the following conditions to be a probability distribution:

- P must map the random variable into probabilities, i.e. $P(x)$ should be a number between 0 and 1.
- $\sum P(x) = 1$, where we sum over all possible values of x . Sometimes rounding errors will cause these sums to be slightly above or below 1. In the continuous case, this summation becomes an **integral**.

Example

Example (Probability of Having x Females in a Biological Family with Three Children)

Note: The following distribution is **not** realistic. Why?

Number of Females (x)	$P(x)$
0	$\frac{1}{6}$
1	$\frac{1}{3}$
2	$\frac{1}{3}$
3	$\frac{1}{6}$

Example

Example (Probability of Having x Females in a Biological Family with Three Children)

Note: The following distribution is **not** realistic. Why?

Number of Females (x)	$P(x)$
0	$\frac{1}{6}$
1	$\frac{1}{3}$
2	$\frac{1}{3}$
3	$\frac{1}{6}$

You can graph a probability distribution of a discrete random variable as a histogram. What would the probability histogram look like for this distribution?

Mean, Variance, and Standard Deviation

Definition (Mean)

$$\mu = \sum_{i=1}^n [x_i \cdot P(x_i)]$$

Mean, Variance, and Standard Deviation

Definition (Mean)

$$\mu = \sum_{i=1}^n [x_i \cdot P(x_i)]$$

The mean is the “expected value” of the random variable x . We write this as $E = \mu$.

Mean, Variance, and Standard Deviation

Definition (Variance)

$$\sigma^2 = \sum_{i=1}^n [(x_i - \mu)^2 \cdot P(x_i)] = \sum_{i=1}^n [x_i^2 \cdot P(x_i)] - \mu^2$$

Mean, Variance, and Standard Deviation

Definition (Variance)

$$\sigma^2 = \sum_{i=1}^n [(x_i - \mu)^2 \cdot P(x_i)] = \sum_{i=1}^n [x_i^2 \cdot P(x_i)] - \mu^2$$

Definition (Standard Deviation)

$$\sigma = \sqrt{\sum_{i=1}^n [x_i^2 \cdot P(x_i)] - \mu^2}$$

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

Number of Females (x)	$P(x)$	$x \cdot P(x)$
0	$\frac{1}{6}$	0
1	$\frac{1}{3}$	0.333
2	$\frac{1}{3}$	0.666
3	$\frac{1}{6}$	0.5
		$\sum x_i \cdot P(x_i)$

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

Number of Females (x)	$P(x)$	$x \cdot P(x)$
0	$\frac{1}{6}$	0
1	$\frac{1}{3}$	0.333
2	$\frac{1}{3}$	0.666
3	$\frac{1}{6}$	0.5
		$\sum x_i \cdot P(x_i) = 1.5 = \mu$

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

# of Females (x)	$P(x)$	x_i^2	$x_i^2 \cdot P(x_i)$
0	$\frac{1}{6}$	0	0
1	$\frac{1}{3}$	1	0.333
2	$\frac{1}{3}$	4	1.333
3	$\frac{1}{6}$	9	1.5
			$\sum x_i^2 \cdot P(x_i) - \mu^2$

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

# of Females (x)	$P(x)$	x_i^2	$x_i^2 \cdot P(x_i)$
0	$\frac{1}{6}$	0	0
1	$\frac{1}{3}$	1	0.333
2	$\frac{1}{3}$	4	1.333
3	$\frac{1}{6}$	9	1.5
			$\sum x_i^2 \cdot P(x_i) - \mu^2 = 0.9166 = \sigma^2$

Mean, Variance, and Standard Deviation: Example

Let's calculate μ , σ^2 , and σ for our previous example.

# of Females (x)	$P(x)$	x_i^2	$x_i^2 \cdot P(x_i)$
0	$\frac{1}{6}$	0	0
1	$\frac{1}{3}$	1	0.333
2	$\frac{1}{3}$	4	1.333
3	$\frac{1}{6}$	9	1.5
			$\sum x_i^2 \cdot P(x_i) - \mu^2 = 0.9166 = \sigma^2$

Thus, $\sigma = 0.9574$.

Unusual Values

We can construct ranges for usual values using these calculated means and standard deviations as well.

Unusual Values

We can construct ranges for usual values using these calculated means and standard deviations as well. We use the same range rule for usual values as before: $[\mu - 2\sigma, \mu + 2\sigma]$.

Unusual Values

We can construct ranges for usual values using these calculated means and standard deviations as well. We use the same range rule for usual values as before: $[\mu - 2\sigma, \mu + 2\sigma]$.

We may also determine if a value is unusually high or low by using the 5% rule:

Unusual Values

We can construct ranges for usual values using these calculated means and standard deviations as well. We use the same range rule for usual values as before: $[\mu - 2\sigma, \mu + 2\sigma]$.

We may also determine if a value is unusually high or low by using the 5% rule:

- A value X is unusually low if $P(x \leq X) \leq 0.05$.

Unusual Values

We can construct ranges for usual values using these calculated means and standard deviations as well. We use the same range rule for usual values as before: $[\mu - 2\sigma, \mu + 2\sigma]$.

We may also determine if a value is unusually high or low by using the 5% rule:

- A value X is unusually low if $P(x \leq X) \leq 0.05$.
- A value X is unusually high if $P(x \geq X) \leq 0.05$.

Binomial Probability Distribution

Definition (Binomial Probability Distribution)

A probability distribution is a **binomial probability distribution** if the following criteria are met:

- The procedure has a fixed number of trials n .

Binomial Probability Distribution

Definition (Binomial Probability Distribution)

A probability distribution is a **binomial probability distribution** if the following criteria are met:

- The procedure has a fixed number of trials n .
- The trials are independent.

Binomial Probability Distribution

Definition (Binomial Probability Distribution)

A probability distribution is a **binomial probability distribution** if the following criteria are met:

- The procedure has a fixed number of trials n .
- The trials are independent.
- The results of the trial must have only two possible outcomes - often called “success” and “failure”.

Binomial Probability Distribution

Definition (Binomial Probability Distribution)

A probability distribution is a **binomial probability distribution** if the following criteria are met:

- The procedure has a fixed number of trials n .
- The trials are independent.
- The results of the trial must have only two possible outcomes - often called “success” and “failure”.
- The probability for success must remain the same throughout all trials.

Binomial Probability Distribution: Example

Let's investigate our example of the number of females in a 3-children family.

Binomial Probability Distribution: Example

Let's investigate our example of the number of females in a 3-children family. First, note that all the requirements to be a binomial probability distribution are met! There are successes (having a female) and failures (having a non-female) as the only outcomes.

Binomial Probability Distribution: Example

Let's investigate our example of the number of females in a 3-children family. First, note that all the requirements to be a binomial probability distribution are met! There are successes (having a female) and failures (having a non-female) as the only outcomes. We let x be our number of successes in n trials (which is fixed at three).

Binomial Probability Distribution: Example

Let's investigate our example of the number of females in a 3-children family. First, note that all the requirements to be a binomial probability distribution are met! There are successes (having a female) and failures (having a non-female) as the only outcomes. We let x be our number of successes in n trials (which is fixed at three). The trials are independent.

Binomial Probability Distribution: Example

Let's investigate our example of the number of females in a 3-children family. First, note that all the requirements to be a binomial probability distribution are met! There are successes (having a female) and failures (having a non-female) as the only outcomes. We let x be our number of successes in n trials (which is fixed at three). The trials are independent. And the probability ($p = 0.5$) of success is the same throughout the whole set of trials.

Binomial Probability Distribution: Example

What's the probability $P(x = 0)$? That would be the probability of having no females: MMM.

Binomial Probability Distribution: Example

What's the probability $P(x = 0)$? That would be the probability of having no females: MMM.

What's the probability $P(x = 1)$? That would be the probability of having one female: FMM, MFM, MMF.

Binomial Probability Distribution: Example

What's the probability $P(x = 0)$? That would be the probability of having no females: MMM.

What's the probability $P(x = 1)$? That would be the probability of having one female: FMM, MFM, MMF.

What's the probability $P(x = 2)$? That would be the probability of having two females: FFM, FMF, MFF.

Binomial Probability Distribution: Example

What's the probability $P(x = 0)$? That would be the probability of having no females: MMM.

What's the probability $P(x = 1)$? That would be the probability of having one female: FMM, MFM, MMF.

What's the probability $P(x = 2)$? That would be the probability of having two females: FFM, FMF, MFF.

What's the probability $P(x = 3)$? That would be the probability of having two females: FFF.

Binomial Probability Distribution: Example

What's the probability $P(x = 0)$? That would be the probability of having no females: MMM.

What's the probability $P(x = 1)$? That would be the probability of having one female: FMM, MFM, MMF.

What's the probability $P(x = 2)$? That would be the probability of having two females: FFM, FMF, MFF.

What's the probability $P(x = 3)$? That would be the probability of having three females: FFF.

So $P(x = 0) = \frac{1}{8}$, $P(x = 1) = \frac{3}{8}$, $P(x = 2) = \frac{3}{8}$, $P(x = 3) = \frac{1}{8}$.

Binomial Probability Distribution: Calculating Probabilities

In a binomial distribution with n trials and probability of success p and failure q , where $p = 1 - q$, you can calculate $P(x = K)$, where K is a whole number less than n in the following way:

Binomial Probability Distribution: Calculating Probabilities

In a binomial distribution with n trials and probability of success p and failure q , where $p = 1 - q$, you can calculate $P(x = K)$, where K is a whole number less than n in the following way:

$$\binom{n}{K} p^K q^{n-K}.$$

Binomial Probability Distribution: Calculating Probabilities

In a binomial distribution with n trials and probability of success p and failure q , where $p = 1 - q$, you can calculate $P(x = K)$, where K is a whole number less than n in the following way:

$$\binom{n}{K} p^K q^{n-K}.$$

Here, $\binom{n}{K}$ (read n choose K) simply denotes $\frac{n!}{(n-K)!K!}$.

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. If you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 5 of them will live in walking distance to school?

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. If you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 5 of them will live in walking distance to school?

Can this even be modeled by a binomial distribution?

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. If you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 5 of them will live in walking distance to school?

Can this even be modeled by a binomial distribution? Strictly speaking, it can't. Why?

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. If you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 5 of them will live in walking distance to school?

Can this even be modeled by a binomial distribution? Strictly speaking, it can't. Why? But we can approximate a selection of 20 students without replacement as independent because the 20 students selected represent less than 5% of the population under consideration.

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. If you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 5 of them will live in walking distance to school?

Can this even be modeled by a binomial distribution? Strictly speaking, it can't. Why? But we can approximate a selection of 20 students without replacement as independent because the 20 students selected represent less than 5% of the population under consideration. So,

$$P(x = 5) = \binom{20}{5} (0.2)^5 (0.8)^{20-5} = 15504 (0.2)^5 (0.8)^{15} \approx 0.175.$$

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. We you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 10 or more of them will live in walking distance to school?

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. We you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 10 or more of them will live in walking distance to school?

This is

$$P(x \geq 10) = \sum_{i=10}^{20} \binom{20}{i} (0.2)^i (0.8)^{20-i} \approx 0.0026.$$

Example

At Arlington High School, 20% of the student body of 1200 students is in walking distance to class. We you randomly select 20 students (without replacement) from the 1200, what is the probability that exactly 10 or more of them will live in walking distance to school?

This is

$$P(x \geq 10) = \sum_{i=10}^{20} \binom{20}{i} (0.2)^i (0.8)^{20-i} \approx 0.0026.$$

In this case, we would say that 10 is an unusually high number of successes.

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4)$

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2)$

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2) \approx 0.156$.

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2) \approx 0.156$.
- $P(x \geq 7)$

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2) \approx 0.156$.
- $P(x \geq 7) \approx 0.057$.

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2) \approx 0.156$.
- $P(x \geq 7) \approx 0.057$.
- $P(x \leq 0)$

Example: Binomial(15,0.25)

Use StatCrunch to determine

- $P(x > 4) \approx 0.314$.
- $P(x = 2) \approx 0.156$.
- $P(x \geq 7) \approx 0.057$.
- $P(x \leq 0) \approx 0.013$.

Parameters for Binomial Distributions

If X is a binomial random variable with n trials and p successes (often written $X \sim \text{Binomial}(n, p)$), then you can calculate mean, variance, and mode using simply n and p .

Parameters for Binomial Distributions

If X is a binomial random variable with n trials and p successes (often written $X \sim \text{Binomial}(n, p)$), then you can calculate mean, variance, and mode using simply n and p .

- $\mu = np$.

Parameters for Binomial Distributions

If X is a binomial random variable with n trials and p successes (often written $X \sim \text{Binomial}(n, p)$), then you can calculate mean, variance, and mode using simply n and p .

- $\mu = np$.
- $\sigma^2 = npq$.

Parameters for Binomial Distributions

If X is a binomial random variable with n trials and p successes (often written $X \sim \text{Binomial}(n, p)$), then you can calculate mean, variance, and mode using simply n and p .

- $\mu = np.$
- $\sigma^2 = npq.$
- $\sigma = \sqrt{npq}$

Parameters for Binomial Distributions

If X is a binomial random variable with n trials and p successes (often written $X \sim \text{Binomial}(n, p)$), then you can calculate mean, variance, and mode using simply n and p .

- $\mu = np$.
- $\sigma^2 = npq$.
- $\sigma = \sqrt{npq}$

With this information, we could use the range rule of thumb rather than the 5% rule of thumb to determine if a value is typical or not.

Example: Binomial(15,0.25)

Using the range rule of thumb, determine if the value $x = 1$ is typical.

Example: Binomial(15,0.25)

Using the range rule of thumb, determine if the value $x = 1$ is typical.

Well, $\mu = 15(0.25) = 3.75$ and $\sigma = \sqrt{15(0.25)(0.75)} \approx 1.677$.

Example: Binomial(15,0.25)

Using the range rule of thumb, determine if the value $x = 1$ is typical.

Well, $\mu = 15(0.25) = 3.75$ and $\sigma = \sqrt{15(0.25)(0.75)} \approx 1.677$. So our typical range of values is

$$[3.75 - 2(1.667), 3.75 + 2(1.667)] = [0.396, 7.104].$$

Example: Binomial(15,0.25)

Using the range rule of thumb, determine if the value $x = 1$ is typical.

Well, $\mu = 15(0.25) = 3.75$ and $\sigma = \sqrt{15(0.25)(0.75)} \approx 1.677$. So our typical range of values is

$$[3.75 - 2(1.667), 3.75 + 2(1.667)] = [0.396, 7.104].$$

Because we know the values only take whole numbers, we say that the range of typical values are the whole numbers from 1 to 7.