

Lecture 7: Chapter 7

C C Moxley

UAB Mathematics

10 October 2016

§7.1 Inferential Statistics

We have previously used descriptive statistics to summarize data in a sample. We will now extend these statistics using inferential methods to make generalizations about the parameters of a population.

§7.1 Inferential Statistics

We have previously used descriptive statistics to summarize data in a sample. We will now extend these statistics using inferential methods to make generalizations about the parameters of a population.

Definition (Inferential Statistics)

The use of sample data to make statements about the statistical/probabilistic characteristics of a population.

§7.2 Estimating a Population Proportion

A **population proportion** describes the percentage (or proportion) of a population which has a certain characteristic. The following are important to estimating population proportions and other population parameters.

§7.2 Estimating a Population Proportion

A **population proportion** describes the percentage (or proportion) of a population which has a certain characteristic. The following are important to estimating population proportions and other population parameters.

Definition (Point Estimate)

A **point estimate** is a single value used to estimate a population parameter.

§7.2 Estimating a Population Proportion

A **population proportion** describes the percentage (or proportion) of a population which has a certain characteristic. The following are important to estimating population proportions and other population parameters.

Definition (Point Estimate)

A **point estimate** is a single value used to estimate a population parameter.

E.g. 35% (or 0.35) is a point estimate.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

Definition (Confidence Level)

A **confidence level** is a probability $(1 - \alpha)$ that describes, in some sense, how sure one is that the true population parameter lies within a given confidence interval.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

Definition (Confidence Level)

A **confidence level** is a probability $(1 - \alpha)$ that describes, in some sense, how sure one is that the true population parameter lies within a given confidence interval.

E.g. 95% (or 0.95) is a confidence level.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

Definition (Confidence Level)

A **confidence level** is a probability $(1 - \alpha)$ that describes, in some sense, how sure one is that the true population parameter lies within a given confidence interval.

E.g. 95% (or 0.95) is a confidence level. In this case, $\alpha = 0.05$.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

Definition (Confidence Level)

A **confidence level** is a probability $(1 - \alpha)$ that describes, in some sense, how sure one is that the true population parameter lies within a given confidence interval.

E.g. 95% (or 0.95) is a confidence level. In this case, $\alpha = 0.05$. Often, you are asked to create the confidence interval given a certain desired confidence level.

§7.2 Estimating a Population Proportion

Definition (Confidence Interval)

A **confidence interval** is an interval of values used to “estimate” the true value of a population parameter. It’s sometimes abbreviated CI.

E.g. The interval $(-1.2, -0.8)$ is a confidence interval.

Definition (Confidence Level)

A **confidence level** is a probability $(1 - \alpha)$ that describes, in some sense, how sure one is that the true population parameter lies within a given confidence interval.

E.g. 95% (or 0.95) is a confidence level. In this case, $\alpha = 0.05$. Often, you are asked to create the confidence interval given a certain desired confidence level. In this chapter, we use confidence intervals for informal hypothesis testing.

§7.2 Estimating a Population Proportion

Definition (Critical Value)

A **critical value** is the number on the boarder which separates typical from atypical values. The number $z_{\frac{\alpha}{2}}$ is a critical value which separates an area of $\frac{\alpha}{2}$ in the right tail of the standard normal distribution.

§7.2 Estimating a Population Proportion

Definition (Critical Value)

A **critical value** is the number on the boarder which separates typical from atypical values. The number $z_{\frac{\alpha}{2}}$ is a critical value which separates an area of $\frac{\alpha}{2}$ in the right tail of the standard normal distribution.

Different distributions must have critical values calculated differently!

§7.2 Estimating a Population Proportion

Definition (Margin of Error)

When data from a simple random sample are used to estimate a population proportion, the **margin of error** which we denote **E**, is the maximum likely difference (with probability $1 - \alpha$) between the observed sample proportion \hat{p} and the true value of the proportion p .

§7.2 Estimating a Population Proportion

Definition (Margin of Error)

When data from a simple random sample are used to estimate a population proportion, the **margin of error** which we denote **E**, is the maximum likely difference (with probability $1 - \alpha$) between the observed sample proportion \hat{p} and the true value of the proportion p .

For proportions, we have

$$|p - \hat{p}| \leq E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}}.$$

§7.2 Estimating a Population Proportion

Definition (Margin of Error)

When data from a simple random sample are used to estimate a population proportion, the **margin of error** which we denote **E**, is the maximum likely difference (with probability $1 - \alpha$) between the observed sample proportion \hat{p} and the true value of the proportion p .

For proportions, we have

$$|p - \hat{p}| \leq E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}}.$$

You can also use this information to calculate how big a sample size needs to be to yield a certain level of confidence with a certain error size. (Always round up when doing this.)

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table. We get $z_{0.025} = 1.96$.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table. We get $z_{0.025} = 1.96$. Then we calculate:

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 1.96 \sqrt{\frac{(0.73)(0.27)}{1007}} \approx 0.027.$$

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table. We get $z_{0.025} = 1.96$. Then we calculate:

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 1.96 \sqrt{\frac{(0.73)(0.27)}{1007}} \approx 0.027.$$

So, our 95% confidence interval is $(0.73 - 0.027, 0.73 + 0.027) = (0.703, 0.757)$.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table. We get $z_{0.025} = 1.96$. Then we calculate:

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 1.96 \sqrt{\frac{(0.73)(0.27)}{1007}} \approx 0.027.$$

So, our 95% confidence interval is $(0.73 - 0.027, 0.73 + 0.027) = (0.703, 0.757)$.

Can we safely say that at least 70% of US adults know what Snapchat is?

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. Construct a 95% confidence interval for the proportion of US adults who know what Snapchat is.

We need to first calculate $z_{0.025}$ using StatCrunch or a table. We get $z_{0.025} = 1.96$. Then we calculate:

$$E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 1.96 \sqrt{\frac{(0.73)(0.27)}{1007}} \approx 0.027.$$

So, our 95% confidence interval is $(0.73 - 0.027, 0.73 + 0.027) = (0.703, 0.757)$.

Can we safely say that at least 70% of US adults know what Snapchat is? Yes!

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. How large should we make the sample in order to be 99% sure that the population proportion lies within 0.02 of the measured sample proportion?

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. How large should we make the sample in order to be 99% sure that the population proportion lies within 0.02 of the measured sample proportion?

We need to first calculate $z_{0.005}$ using StatCrunch or a table.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. How large should we make the sample in order to be 99% sure that the population proportion lies within 0.02 of the measured sample proportion?

We need to first calculate $z_{0.005}$ using StatCrunch or a table. We get $z_{0.005} = 2.575$.

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. How large should we make the sample in order to be 99% sure that the population proportion lies within 0.02 of the measured sample proportion?

We need to first calculate $z_{0.005}$ using StatCrunch or a table. We get $z_{0.005} = 2.575$. Then we use the formula:

$$0.02 = E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 2.576 \sqrt{\frac{(0.73)(0.27)}{n}},$$

§7.2 Example

Pew Research polled 1007 randomly selected US adults and showed that 73% of the respondents knew what Snapchat was. How large should we make the sample in order to be 99% sure that the population proportion lies within 0.02 of the measured sample proportion?

We need to first calculate $z_{0.005}$ using StatCrunch or a table. We get $z_{0.005} = 2.575$. Then we use the formula:

$$0.02 = E = z_{\frac{\alpha}{2}} \sqrt{\frac{\hat{p}\hat{q}}{n}} = 2.576 \sqrt{\frac{(0.73)(0.27)}{n}},$$

which implies that $n \approx 3269.78$, so we need a sample size of at least 3270.

§7.2 Requirements for Using Proportion Estimation Methods

Note: The methods used in the previous examples are only appropriate when

§7.2 Requirements for Using Proportion Estimation Methods

Note: The methods used in the previous examples are only appropriate when

- The sample is a simple random sample.

§7.2 Requirements for Using Proportion Estimation Methods

Note: The methods used in the previous examples are only appropriate when

- The sample is a simple random sample.
- The conditions for a binomial distribution are satisfied.

§7.2 Requirements for Using Proportion Estimation Methods

Note: The methods used in the previous examples are only appropriate when

- The sample is a simple random sample.
- The conditions for a binomial distribution are satisfied.
- There are at least five successes and five failures.

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or
- The sample must be of size $n > 30$.

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or
- The sample must be of size $n > 30$.

If σ is known, we can just apply the Central Limit Theorem to obtain a proper confidence interval, using

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or
- The sample must be of size $n > 30$.

If σ is known, we can just apply the Central Limit Theorem to obtain a proper confidence interval, using

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}.$$

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or
- The sample must be of size $n > 30$.

If σ is known, we can just apply the Central Limit Theorem to obtain a proper confidence interval, using

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}.$$

If σ is not known, we must use the Student t distribution rather than the standard normal to calculate our critical value:

§7.3 Estimating a Population Mean

We use the same notions as in §6.2, but we do have different requirements:

- The population must have a normal distribution or
- The sample must be of size $n > 30$.

If σ is known, we can just apply the Central Limit Theorem to obtain a proper confidence interval, using

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}.$$

If σ is not known, we must use the Student t distribution rather than the standard normal to calculate our critical value:

$$E = t_{\frac{\alpha}{2}} \frac{s}{\sqrt{n}}.$$

§7.3 Estimating a Population Mean

Note: If you need to calculate $t_{\frac{\alpha}{2}}$, you need to know how many **degrees of freedom** the t distribution has.

§7.3 Estimating a Population Mean

Note: If you need to calculate $t_{\frac{\alpha}{2}}$, you need to know how many **degrees of freedom** the t distribution has. The degrees of freedom of t is always given by

$$n - 1,$$

where n is the sample size.

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$. It's known that cats have body temperatures which have a standard deviation of $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$. It's known that cats have body temperatures which have a standard deviation of $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$. It's known that cats have body temperatures which have a standard deviation of $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} = 1.96 \frac{0.7}{\sqrt{50}} \approx 0.19,$$

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$. It's known that cats have body temperatures which have a standard deviation of $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} = 1.96 \frac{0.7}{\sqrt{50}} \approx 0.19,$$

so our 95% confidence interval is $101.5 \pm 0.19^{\circ}F$ or

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$. It's known that cats have body temperatures which have a standard deviation of $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} = 1.96 \frac{0.7}{\sqrt{50}} \approx 0.19,$$

so our 95% confidence interval is $101.5 \pm 0.19^{\circ}F$ or $(101.31, 101.69)$.

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$ and their standard deviation was $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$ and their standard deviation was $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$ and their standard deviation was $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = t_{\frac{\alpha}{2}} \frac{s}{\sqrt{n}} = 2.0096 \frac{0.7}{\sqrt{50}} \approx 0.20,$$

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$ and their standard deviation was $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = t_{\frac{\alpha}{2}} \frac{s}{\sqrt{n}} = 2.0096 \frac{0.7}{\sqrt{50}} \approx 0.20,$$

so our 95% confidence interval is $101.5 \pm 0.20^{\circ}F$ or

§7.3 Example

50 body temperatures of cats were taken and their average was $101.5^{\circ}F$ and their standard deviation was $0.7^{\circ}F$. Find a 95% confidence interval for the mean body temperature of cats.

We need to calculate

$$E = t_{\frac{\alpha}{2}} \frac{s}{\sqrt{n}} = 2.0096 \frac{0.7}{\sqrt{50}} \approx 0.20,$$

so our 95% confidence interval is $101.5 \pm 0.20^{\circ}F$ or $(101.30, 101.70)$.

§7.3 Example

Assuming that the standard deviation of cats' body temperatures is known to be $0.7^{\circ}F$. How many cats would you need to sample to get a 90% confidence interval with error of $0.1^{\circ}F$?

§7.3 Example

Assuming that the standard deviation of cats' body temperatures is known to be $0.7^{\circ}F$. How many cats would you need to sample to get a 90% confidence interval with error of $0.1^{\circ}F$?

We solve for n the equation

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \implies$$

§7.3 Example

Assuming that the standard deviation of cats' body temperatures is known to be $0.7^{\circ}F$. How many cats would you need to sample to get a 90% confidence interval with error of $0.1^{\circ}F$?

We solve for n the equation

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \implies 0.1 = 1.645 \frac{0.7}{\sqrt{n}}.$$

§7.3 Example

Assuming that the standard deviation of cats' body temperatures is known to be $0.7^{\circ}F$. How many cats would you need to sample to get a 90% confidence interval with error of $0.1^{\circ}F$?

We solve for n the equation

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \implies 0.1 = 1.645 \frac{0.7}{\sqrt{n}}.$$

This yields

$$n = \left(\frac{z_{\frac{\alpha}{2}} \sigma}{E} \right)^2 = \left(\frac{1.645(0.7)}{0.1} \right)^2 \approx 132.6,$$

§7.3 Example

Assuming that the standard deviation of cats' body temperatures is known to be $0.7^{\circ}F$. How many cats would you need to sample to get a 90% confidence interval with error of $0.1^{\circ}F$?

We solve for n the equation

$$E = z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}} \implies 0.1 = 1.645 \frac{0.7}{\sqrt{n}}.$$

This yields

$$n = \left(\frac{z_{\frac{\alpha}{2}} \sigma}{E} \right)^2 = \left(\frac{1.645(0.7)}{0.1} \right)^2 \approx 132.6,$$

which means our sample size must be at least 133 cats.

§7.4 Estimating Population Standard Deviation or Variance

As before, we still have the notions of point estimate and confidence interval, but we must use the **chi-squared distribution**, denoted χ^2 , in order to calculate the critical value!

§7.4 Estimating Population Standard Deviation or Variance

As before, we still have the notions of point estimate and confidence interval, but we must use the **chi-squared distribution**, denoted χ^2 , in order to calculate the critical value!

Just as with the Student t distribution, you must know the degrees of freedom of a χ^2 distribution, which is again given by

$$n - 1.$$

Note:

§7.4 Estimating Population Standard Deviation or Variance

As before, we still have the notions of point estimate and confidence interval, but we must use the **chi-squared distribution**, denoted χ^2 , in order to calculate the critical value!

Just as with the Student t distribution, you must know the degrees of freedom of a χ^2 distribution, which is again given by

$$n - 1.$$

Note: The χ^2 distribution is **not symmetric**, and neither is the confidence interval it creates.

§7.4 Estimating Population Standard Deviation or Variance

As before, we still have the notions of point estimate and confidence interval, but we must use the **chi-squared distribution**, denoted χ^2 , in order to calculate the critical value!

Just as with the Student t distribution, you must know the degrees of freedom of a χ^2 distribution, which is again given by

$$n - 1.$$

Note: The χ^2 distribution is **not symmetric**, and neither is the confidence interval it creates. For this reason, we never denote a confidence interval for the standard deviation in the $\sigma \pm E$ notation!

§7.4 Estimating Population Standard Deviation or Variance


As before, we still have the notions of point estimate and confidence interval, but we must use the **chi-squared distribution**, denoted χ^2 , in order to calculate the critical value!

Just as with the Student t distribution, you must know the degrees of freedom of a χ^2 distribution, which is again given by

$$n - 1.$$

Note: The χ^2 distribution is **not symmetric**, and neither is the confidence interval it creates. For this reason, we never denote a confidence interval for the standard deviation in the $\sigma \pm E$ notation! We always use interval notation!

§7.4 The χ^2 Distribution



§7.4 Estimating Population Standard Deviation or Variance

For the standard deviation, we have

$$\sqrt{\frac{(n-1)s^2}{\chi_R^2}} < \sigma < \sqrt{\frac{(n-1)s^2}{\chi_L^2}},$$

§7.4 Estimating Population Standard Deviation or Variance

For the standard deviation, we have

$$\sqrt{\frac{(n-1)s^2}{\chi_R^2}} < \sigma < \sqrt{\frac{(n-1)s^2}{\chi_L^2}},$$

and for the variance, we have

$$\frac{(n-1)s^2}{\chi_R^2} < \sigma^2 < \frac{(n-1)s^2}{\chi_L^2}.$$

§7.4 Estimating Population Standard Deviation or Variance

We have the following requirements when estimating the standard deviation:

§7.4 Estimating Population Standard Deviation or Variance

We have the following requirements when estimating the standard deviation:

- The sample must be a simple random sample.

§7.4 Estimating Population Standard Deviation or Variance

We have the following requirements when estimating the standard deviation:

- The sample must be a simple random sample.
- The population **must be normally distributed, even if the sample is large!**

§7.4 Example

The time a student spends on a section of the SAT is given in minutes: 25, 21, 21, 18, 13, 13, 14, 16, 12, 22, 25, 25, 30. These times are known to be normally distributed. Calculate a 90% confidence interval for the standard deviation of these times.

§7.4 Example

The time a student spends on a section of the SAT is given in minutes: 25, 21, 21, 18, 13, 13, 14, 16, 12, 22, 25, 25, 30. These times are known to be normally distributed. Calculate a 90% confidence interval for the standard deviation of these times.

We have that $\bar{x} = 19.62$ and $s = 5.752$.

§7.4 Example

The time a student spends on a section of the SAT is given in minutes: 25, 21, 21, 18, 13, 13, 14, 16, 12, 22, 25, 25, 30. These times are known to be normally distributed. Calculate a 90% confidence interval for the standard deviation of these times.

We have that $\bar{x} = 19.62$ and $s = 5.752$. (We get this from StatCruch, using Summary Stats by column).

§7.4 Example

The time a student spends on a section of the SAT is given in minutes: 25, 21, 21, 18, 13, 13, 14, 16, 12, 22, 25, 25, 30. These times are known to be normally distributed. Calculate a 90% confidence interval for the standard deviation of these times.

We have that $\bar{x} = 19.62$ and $s = 5.752$. (We get this from StatCruch, using Summary Stats by column).

Therefore,

$$\sqrt{\frac{(n-1)s^2}{\chi_R^2}} < \sigma < \sqrt{\frac{(n-1)s^2}{\chi_L^2}} \implies$$
$$4.35 = \sqrt{\frac{(12)5.752^2}{21.0261}} < \sigma < \sqrt{\frac{(12)5.752^2}{5.2260}} = 8.72.$$

§7.4 Example

So we have that the 90% confidence interval is (4.35,8.68).

§7.4 Example

So we have that the 90% confidence interval is $(4.35, 8.68)$. What does this actually mean?

§7.4 Example

So we have that the 90% confidence interval is $(4.35, 8.68)$. What does this actually mean?

This means that if we took a large number of simple random samples of size 13 and calculated their standard deviation, then 90% of the time, your calculated standard deviation would fall into the interval $(4.35, 8.68)$.

§7.4 Example

So we have that the 90% confidence interval is (4.35,8.68). What does this actually mean?

This means that if we took a large number of simple random samples of size 13 and calculated their standard deviation, then 90% of the time, your calculated standard deviation would fall into the interval (4.35,8.68). It does not mean that there is a 90% chance that σ falls within the interval.

§7.4 Precise Language

What does it mean to have a 95% CI for population mean?

§7.4 Precise Language

What does it mean to have a 95% CI for population mean? It is precisely an interval which, when a random selection is made from the sampling distribution of the mean, will contain the given selection 95% of the time. We'll talk about how this relates to hypothesis testing in the next chapter, and it will be much easier to speak about what a 95% CI (or confidence level) means in the context of hypothesis testing.

In a probability histogram for the number of female girls in a family of 5, how would you determine the probability of having two or three females?

In some population, the mean is 25 and the standard deviation is 6. What is the probability that, in a simple random sample of 36, the sample mean would be between 24 and 26?

20% of students at UAB live on campus. Approximate the probability of choosing 10 students at random from the student population and having all of them live on campus.

What is the probability of having exactly one student who does not live on campus under the same circumstances as in the previous question?

What is the z-score of a test with a score of 75 if the mean on this test is 80 with a standard deviation of 2.5?

Describe in words what is meant by the sampling distribution of a median.

§Review

In a bag, you have 10 coins: 5 quarters, 1 dime, 1 nickel, and 3 pennies. If you pay 15 cents, you can draw one coin from the bag at random. Is this game a good game to pay many times?

The Central Limit Theorem says that when a sample is large the distribution of a sample mean is always what?

What's the probability of selecting two kings from a deck of cards with and without replacement? When does it make sense to compute probabilities when sampling without replacement as if we are sampling with replacement?

How many ways are there of getting two heads when you're flipping a coin 5 times?

Make sure to know all theorems, rules, and laws!