

Rotation Sets of Billiards with One Obstacle

Alexander Blokh^{1,*}, Michał Misiurewicz^{2,**}, Nándor Simányi^{1,***}

¹ Department of Mathematics, University of Alabama at Birmingham, University Station, Birmingham, AL 35294-2060, USA. E-mail: ablokh@math.uab.edu; simanyi@math.uab.edu

² Department of Mathematical Sciences, IUPUI, 402 N. Blackford Street, Indianapolis, IN 46202-3216, USA. E-mail: mmisiure@math.iupui.edu

Received: 16 August 2005 / Accepted: 10 December 2005
Published online: 14 April 2006 – © Springer-Verlag 2006

Abstract: We investigate the rotation sets of billiards on the m -dimensional torus with one small convex obstacle and in the square with one small convex obstacle. In the first case the displacement function, whose averages we consider, measures the change of the position of a point in the universal covering of the torus (that is, in the Euclidean space), in the second case it measures the rotation around the obstacle. A substantial part of the rotation set has usual strong properties of rotation sets.

1. Introduction

Traditionally, billiards have been investigated from the point of view of Ergodic Theory. That is, the properties that have been studied, were the statistical properties with respect to the natural invariant measure equivalent to the Lebesgue measure. However, it is equally important to investigate the limit behavior of *all* trajectories and not only of *almost all* of them. In particular, periodic trajectories (which are of zero measure) are of great interest. A widely used method in this context is to observe that billiards in convex domains are twist maps (see, e.g. [6], Sect. 9.2), so the well developed rotation theory for twist maps (see, e.g. [6], Sect. 9.3) applies to them.

Rotation Theory has been recently developed further and its scope has been significantly widened (see, e.g., Chapter 6 of [1] for a brief overview). This opens possibilities of its application to new classes of billiards. In the general Rotation Theory one considers a dynamical system together with an *observable*, that is a function on the phase space, with values in a vector space. Then one takes limits of ergodic averages of the observable along longer and longer pieces of trajectories. The *rotation set* obtained in such a way contains all averages of the observable along periodic orbits, and, by the

* The first author was partially supported by NSF grant DMS 0456748.

** The second author was partially supported by NSF grant DMS 0456526.

*** The third author was partially supported by NSF grant DMS 0457168.

Birkhoff Ergodic Theorem, integrals of the observable with respect to all ergodic invariant probability measures. With the natural choice of an observable, information about the appropriate rotation set allows one to describe the behavior of the trajectories of the system (see examples in Chapter 6 of [1]). Exact definitions are given later in the paper.

We have to stress again that we consider *all* trajectories. Indeed, restricting attention to one ergodic measure would result in seeing only one rotation vector. However, rotation vectors of other points, non-typical for the measure, will be missing. Thus, the approach to the billiards should be from the point of view of Topological Dynamics, instead of Ergodic Theory, even though we do consider various invariant measures for which rotation vector can be computed. Observe that the rotation set for a suitably chosen observable is a useful characteristic of the dynamical system.

In the simplest case, the observable is the increment in one step (for discrete systems) or the derivative (for systems with continuous time) of another function, called *displacement*. When the displacement is chosen in a natural way, the results on the rotation set are especially interesting. In this paper we consider two similar classes of billiards, and the observables which we use for them are exactly of that type. One system consists of billiards on an m -dimensional torus with one small convex obstacle which we lift to the universal covering of the torus (that is, to the Euclidean space) and consider the natural displacement there. Those models constitute the rigorous mathematical formulation of the so called Lorentz gas dynamics with periodic configuration of obstacles. They are especially important for physicists doing research in the foundations of nonequilibrium dynamics, since the Lorentz gas serves as a good paradigm for nonequilibrium stationary states, see the nice survey [5].

The other system consists of billiards in a square with one small convex obstacle close to the center of the square; here we measure average rotation around a chosen obstacle using the argument as the displacement.

We treat both billiards as flows. This is caused by the fact that in the lifting (or unfolding for a billiard in a square) we may have infinite horizon, especially if the obstacle is small. In other words, there are infinite trajectories without reflections, so when considering billiards as maps, we would have to divide by zero. Although this is not so bad by itself (infinity exists), we lose compactness and cannot apply nice general machinery of the rotation theory (see, e.g., [11]).

Note that in the case of a billiard in a square we have to deal with trajectories that reflect from the vertices of the square. We can think about such reflection as two infinitesimally close reflections from two adjacent sides. Then it is clear that our trajectory simply comes back along the same line on which it arrived at the vertex, and that this does not destroy the continuity of the flow.

The ideas, methods, and results in both cases, the torus and the square, are very similar. However, there are some important differences, and, in spite of its two-dimensionality, in general the square case is more complicated. Therefore we decided to treat the torus case first (in Sects. 2, 3 and 4) and then, when we describe the square case (in Sect. 5), we describe the differences from the torus case, without repeating the whole proofs. We believe that this type of exposition is simpler for a reader than the one that treats both cases simultaneously or the one that produces complicated abstract theorems that are then applied in both cases. In Sect. 6 we get additionally some general results, applicable also to other situations.

Let us describe briefly the main results of the paper. The exact definitions will be given later. Let us only note that the *admissible rotation set* is a subset of the full rotation set, about which we can prove much stronger results than about the full rotation

set. Also, a *small* obstacle does not mean an “arbitrarily small” one. We derive various estimates of the size of the admissible rotation set. In the torus case the estimates that are independent of the dimension are non-trivial because of the behavior of the geometry of \mathbb{R}^m as $m \rightarrow \infty$. In both cases we show that the admissible rotation set approximates better and better the full rotation set when the size of the obstacle diminishes.

We prove that in both cases, the torus and the square, if the obstacle is small, then the admissible rotation set is convex, rotation vectors of periodic orbits are dense in it, and if u is a vector from its interior, then there exists a trajectory with the rotation vector u (and even an ergodic invariant measure, for which the integral of the velocity is equal to u , so that u is the rotation vector of almost every trajectory). The full rotation set is connected, and in the case of the square, is equal to the interval $[-\sqrt{2}/4, \sqrt{2}/4]$.

We conjecture that the full rotation set shares the strong properties of the admissible rotation set.

2. Preliminary Results - Torus

Let us consider a billiard on the m -dimensional torus $\mathbb{T}^m = \mathbb{R}^m / \mathbb{Z}^m$ ($m \geq 2$) with one strictly convex (that is, it is convex and its boundary does not contain any straight line segment) obstacle O with a smooth boundary. We do not specify explicitly how large the obstacle is, but let us think about it as a rather small one. When we lift the whole picture to \mathbb{R}^m then we get a family of obstacles O_k , where $k \in \mathbb{Z}^m$ and O_k is O_0 translated by the vector k .

When we speak about a *trajectory*, we mean a positive (one-sided) billiard trajectory, unless we explicitly say that it is a full (two-sided) one. However, we may mean a trajectory in the phase space, in the configuration space (on the torus), or in the lifting or unfolding (the Euclidean space). It will be usually clear from the context, which case we consider.

We will say that the obstacle O_k is *between* O_i and O_j if it intersects the convex hull of $O_i \cup O_j$ and $k \neq i, j$. For a trajectory P , beginning on a boundary of O , its *type* is a sequence $(k_n)_{n=0}^\infty$ of elements of \mathbb{Z}^m if the continuous lifting of P to \mathbb{R}^m that starts at the boundary of O_{k_0} reflects consecutively from O_{k_n} , $n = 1, 2, \dots$. In order to make the type unique for a given P , we will additionally assume that $k_0 = \mathbf{0}$. Note that except for the case when P at its initial point is tangent to O , there are infinitely many reflections, so the type of P is well defined. This follows from the following lemma. In it, we do not count tangency as a reflection.

Lemma 2.1. *If a trajectory has one reflection then it has infinitely many reflections.*

Proof. Suppose that a trajectory has a reflection, but there are only finitely many of them. Then we can start the trajectory from the last reflection. Its ω -limit set is an affine subtorus of \mathbb{T}^m and the whole (positive) trajectory is contained in this subtorus (and it is dense there). Since we started with a reflection, this subtorus intersects the interior of the obstacle. Since the trajectory is dense in this subtorus, we get a contradiction. \square

Of course, there may be trajectories without any reflections. In particular, if the obstacle is contained in a ball of radius less than $1/2$, there are such trajectories in the direction of the basic unit vectors.

Sometimes we will speak about the *type* of a piece of a trajectory; then it is a finite sequence. We will also use the term *itinerary*.

We will call a sequence $(k_n)_{n=0}^\infty$ of elements of \mathbb{Z}^m *admissible* if

1. $\mathbf{k}_0 = \mathbf{0}$,
2. for every n we have $\mathbf{k}_{n+1} \neq \mathbf{k}_n$,
3. for every n there is no obstacle between $O_{\mathbf{k}_n}$ and $O_{\mathbf{k}_{n+1}}$,
4. for every n the obstacle $O_{\mathbf{k}_{n+1}}$ is not between $O_{\mathbf{k}_n}$ and $O_{\mathbf{k}_{n+2}}$.

Theorem 2.2. *For any admissible sequence $(\mathbf{k}_n)_{n=0}^\infty$ there is a trajectory with type $(\mathbf{k}_n)_{n=0}^\infty$. If additionally there is $\mathbf{p} \in \mathbb{Z}^m$ and a positive integer q such that $\mathbf{k}_{n+q} = \mathbf{k}_n + \mathbf{p}$ for every n then this trajectory can be chosen periodic of discrete period q (that is, after q reflections we come back to the starting point in the phase space). Similarly, for any admissible sequence $(\mathbf{k}_n)_{n=-\infty}^\infty$ there is a trajectory with type $(\mathbf{k}_n)_{n=-\infty}^\infty$.*

Proof. Fix n . For every sequence $A = (\mathbf{x}_i)_{i=0}^n$ such that \mathbf{x}_i belongs to the boundary of $O_{\mathbf{k}_i}$ for $i = 0, 1, \dots, n$, let $\Gamma(A)$ be the curve obtained by joining consecutive points \mathbf{x}_i by straight segments (such a curve may intersect interiors of some obstacles). Since the Cartesian product of the boundaries of $O_{\mathbf{k}_i}$ is compact and the length of $\Gamma(A)$ depends continuously on A , there is an A for which this length is minimal.

We claim that in such a case $\Gamma(A)$ is a piece of a trajectory. By (3), the segment I_i joining \mathbf{x}_i with \mathbf{x}_{i+1} cannot intersect any obstacle except $O_{\mathbf{k}_i}$ and $O_{\mathbf{k}_{i+1}}$. If it intersects $O_{\mathbf{k}_i}$ at more than one point, it intersects its boundary at \mathbf{x}_i and at another point \mathbf{y} . Then replacing \mathbf{x}_i by \mathbf{y} will make $\Gamma(A)$ shorter, a contradiction. This argument does not work only if \mathbf{x}_{i-1} , \mathbf{x}_i and \mathbf{x}_{i+1} are collinear and \mathbf{x}_i lies between \mathbf{x}_{i-1} and \mathbf{x}_{i+1} . However, such situation is excluded by (4). This proves that I_i does not intersect $O_{\mathbf{k}_i}$ at more than one point. Similarly, it does not intersect $O_{\mathbf{k}_{i+1}}$ at more than one point. Now the known property of curves with minimal lengths guarantees that at every $\mathbf{x}_i, i = 1, 2, \dots, n - 1$, the incidence and reflection angles are equal. This proves our claim. For a two-sided sequence $(\mathbf{k}_n)_{n=-\infty}^\infty$ the argument is very similar.

Now we make this construction for every n and get a sequence $(A_n)_{n=1}^\infty$ of pieces of trajectories. We note their initial points in the phase space (points and directions) and choose a convergent subsequence of those. Then the trajectory of this limit point in the phase space will have the prescribed type.

If there is $\mathbf{p} \in \mathbb{Z}^m$ and a positive integer q such that $\mathbf{k}_{n+q} = \mathbf{k}_n + \mathbf{p}$ for every n , then we consider only the sequence $A = (\mathbf{x}_i)_{i=0}^{q-1}$ and repeat the first part of the above proof adding the segment joining \mathbf{x}_{q-1} with $\mathbf{x}_0 + \mathbf{p}$ to $\Gamma(A)$. \square

Note that by Corollary 1.2 of [4], if the obstacle is strictly convex then a periodic orbit from the above theorem is unique.

The next lemma essentially expresses the fact that any billiard flow with convex obstacles lacks focal points. It follows from the corollary after Lemma 2 of [9]. The types of trajectory pieces about which we speak in this lemma are not necessarily admissible.

Lemma 2.3. *For a given finite sequence $B = (\mathbf{k}_n)_{n=0}^s$ of elements of \mathbb{Z}^m and points $\mathbf{x}_0, \mathbf{x}_s$ on the boundaries of $O_{\mathbf{k}_0}$ and $O_{\mathbf{k}_s}$ respectively, there is at most one trajectory piece of type B starting at \mathbf{x}_0 and ending at \mathbf{x}_s . The same remains true if we allow the first segment of the trajectory piece to cross $O_{\mathbf{k}_0}$ and the last one to cross $O_{\mathbf{k}_s}$ (as in Fig. 2.1).*

Corollary 2.4. *If the trajectory piece from Lemma 2.3 exists and has admissible type, then it is the shortest path of type B starting at \mathbf{x}_0 and ending at \mathbf{x}_s .*

Proof. Similarly as in the proof of Theorem 2.2, the shortest path of type B from \mathbf{x}_0 to \mathbf{x}_s is a trajectory piece (here we allow the first segment of the trajectory piece to cross




Fig. 2.1. A trajectory piece crossing the first and last obstacles

O_{k_0} and the last one to cross O_{k_s}). By Lemma 2.3, it is equal to the trajectory piece from that lemma. \square

Of course this trajectory piece depends on x_0 and x_s . However, its length depends on those two points only up to an additive constant. Denote by c the diameter of O .

Lemma 2.5. *For every admissible finite sequence $B = (k_n)_{n=0}^s$ of elements of \mathbb{Z}^m the lengths of trajectory pieces of type B (even if we allow them to cross O_{k_0} and O_{k_s}) differ by at most $2c$. The displacements along those trajectory pieces also differ by at most $2c$.*

Proof. Let Γ and Γ' be two such trajectory pieces, joining x_0 with x_s and y_0 with y_s respectively, where x_0, y_0 belong to the boundary of O_{k_0} and x_s, y_s belong to the boundary of O_{k_s} . Replace the first segment of Γ by adding to it the segment joining x_0 with y_0 , and do similarly with the last segment of Γ . Then we get a path joining y_0 with y_s of type B . By Corollary 2.4, its length is not smaller than the length of Γ' . On the other hand, its length is not larger than the length of Γ plus $2c$. Performing the same construction with the roles of Γ and Γ' reversed, we conclude that the difference of the lengths of those two paths is not larger than $2c$.

The second statement of the lemma is obvious. \square

One can look at the definition of an admissible sequence in the following way. Instead of a sequence $(k_n)_{n=0}^\infty$ of elements of \mathbb{Z}^m we consider the sequence $(l_n)_{n=1}^\infty$, where $l_n = k_n - k_{n-1}$. Since $k_0 = \mathbf{0}$, knowing $(l_n)_{n=1}^\infty$ we can recover $(k_n)_{n=0}^\infty$. Now, condition (3) can be restated as no obstacle between O_0 and O_{l_n} , and condition (4) as the obstacle O_{l_n} not between O_0 and $O_{l_n+l_{n+1}}$. Let G be the directed graph whose vertices are those $j \in \mathbb{Z}^m \setminus \{\mathbf{0}\}$ for which there is no obstacle between O_0 and O_j , and there is an edge (arrow) from j to i if and only if O_j is not between O_0 and O_{j+i} . Then every sequence $(l_n)_{n=1}^\infty$ obtained from an admissible sequence is a one-sided infinite path in G , and vice versa, each one-sided infinite path in G is a sequence $(l_n)_{n=1}^\infty$ obtained from an admissible sequence. Hence, we can speak about paths *corresponding* to admissible sequences and admissible sequences *corresponding* to paths.

Lemma 2.6. *The set of vertices of G is finite.*

Proof. Fix an interior point x of O_0 . By Lemma 2.1, any ray beginning at x intersects the interior of some O_k with $k \neq \mathbf{0}$. Let V_k be the set of directions (points of the unit sphere) for which the corresponding ray intersects the interior of O_k . This set is open, so we get an open cover of a compact unit sphere. It has a finite subcover, so there exists

a constant $M > 0$ such that every ray from \mathbf{x} of length M intersects the interior of some $O_{\mathbf{k}}$ with $\mathbf{k} \neq \mathbf{0}$. This proves that the set of vertices of G is finite. \square

Note that in G there is never an edge from a vertex to itself. Moreover, there is a kind of symmetry in G . Namely, if \mathbf{k} is a vertex then $-\mathbf{k}$ is a vertex; there is an edge from \mathbf{k} to $-\mathbf{k}$; and if there is an edge from \mathbf{k} to \mathbf{j} then there is an edge from $-\mathbf{j}$ to $-\mathbf{k}$.

The following lemma establishes another symmetry in G .

Lemma 2.7. *If $\mathbf{k}, \mathbf{j} \in \mathbb{Z}^m$ and $O_{\mathbf{k}}$ is between $O_{\mathbf{0}}$ and $O_{\mathbf{k}+\mathbf{j}}$, then $O_{\mathbf{j}}$ is also between $O_{\mathbf{0}}$ and $O_{\mathbf{k}+\mathbf{j}}$. Thus, if there is an edge in G from \mathbf{k} to \mathbf{j} then there is an edge from \mathbf{j} to \mathbf{k} .*

Proof. The map $f(x) = \mathbf{k} + \mathbf{j} - x$ defines an isometry of \mathbb{Z}^m and $f(O_{\mathbf{0}}) = O_{\mathbf{k}+\mathbf{j}}$, $f(O_{\mathbf{k}+\mathbf{j}}) = O_{\mathbf{0}}$, $f(O_{\mathbf{k}}) = O_{\mathbf{j}}$. This proves the first statement of the lemma. The second statement follows from the first one and from the definition of edges in G . \square

We will say that the obstacle O is *small* if it is contained in a closed ball of radius smaller than $\sqrt{2}/4$. To simplify the notation, in the rest of the paper, whenever the obstacle is small, we will be using the lifting to \mathbb{R}^m such that the centers of the balls of radii smaller than $\sqrt{2}/4$ containing the obstacles will be at the points of \mathbb{Z}^m .

Denote by U the set of unit vectors from \mathbb{Z}^m (that is, the ones with one component ± 1 and the rest of components 0), and by A_m the set $\{-1, 0, 1\}^m \setminus \{\mathbf{0}\}$ (we use the subscript m by A , since this set will be used sometimes when we consider all dimensions at once). In particular, $U \subset A_m$.

Lemma 2.8. *Let O be small. If $\mathbf{k}, \mathbf{l} \in \mathbb{Z}^m \setminus \{\mathbf{0}\}$ and $\langle \mathbf{k}, \mathbf{l} \rangle \leq 0$, then $O_{\mathbf{k}}$ is not between $O_{\mathbf{0}}$ and $O_{\mathbf{k}+\mathbf{l}}$. In particular, if \mathbf{k} and \mathbf{l} are vertices of G and $\langle \mathbf{k}, \mathbf{l} \rangle \leq 0$, then there are edges in G from \mathbf{k} to \mathbf{l} and from \mathbf{l} to \mathbf{k} .*

Proof. We will use elementary geometry. Consider the triangle with vertices $A = \mathbf{0}$, $B = \mathbf{k}$ and $C = \mathbf{k} + \mathbf{l}$. The angle at the vertex B is at most $\pi/2$, and the lengths of the sides AB and BC are at least 1. We need to construct a straight line which separates the plane P in which the triangle ABC lies into two half-planes with the first one containing the open disk of radius $\sqrt{2}/4$ centered at B and the second one containing such disks centered at A and C . Then the hyperplane of dimension $m - 1$ through this line and perpendicular to the plane P will separate $O_{\mathbf{k}}$ from $O_{\mathbf{0}}$ and $O_{\mathbf{k}+\mathbf{l}}$. This will prove that there is an edge in G from \mathbf{k} to \mathbf{l} . By Lemma 2.7, there will be also an edge in G from \mathbf{l} to \mathbf{k} .

Let D and E be the points on the sides BA and BC respectively, whose distance from B is $1/2$ and let L be the straight line through D and E . Since the angle at the vertex B is at most $\pi/2$, the distance of B from L is at least $\sqrt{2}/4$. Since $|AD| \geq |BD|$ and $|CE| \geq |BE|$, the distances of A and C from L are at least as large as the distance of B from L . This completes the proof. \square

Lemma 2.9. *For a billiard on a torus with a small obstacle, all elements of A_m are vertices of G .*

Proof. Let $\mathbf{u} \in A_m$ and $\mathbf{v} \in \mathbb{Z}^m \setminus \{\mathbf{0}, \mathbf{u}\}$. If $\mathbf{v} = (v_1, v_2, \dots, v_m)$ then $|\langle \mathbf{v}, \mathbf{u} \rangle| \leq \sum_{i=1}^m |v_i| \leq \|\mathbf{v}\|^2$, so $\langle \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle \leq 0$. Therefore, by Lemma 2.8, \mathbf{v} is not between $\mathbf{0}$ and \mathbf{u} . This proves that \mathbf{u} is a vertex of G . \square

Lemma 2.10. *Assume that O is small. Then G is connected, and for every vertices \mathbf{k}, \mathbf{l} of G there is a path of length at most 3 from \mathbf{k} to \mathbf{l} in G , via elements of U .*

Proof. By Lemma 2.9, the set of vertices of G contains U . Let k, l be vertices of G . Then $k, l \neq \mathbf{0}$, so there exist elements u, v of U such that $\langle k, u \rangle \leq 0$ and $\langle l, v \rangle \leq 0$. By Lemma 2.8, there are edges in G from k to u and from v to l . If $u = v$ then kul is a path of length 2 from k to l . If $u \neq v$ then $\langle u, v \rangle = 0$, so by Lemma 2.8 there is an edge from u to v . Then $kuvl$ is a path of length 3 from k to l . \square

3. Rotation Set - Torus

Now we have enough information in order to start investigating the *rotation set* R of our billiard. It consists of limits of the sequences $((y_n - x_n)/t_n)_{n=1}^\infty$, where there is a trajectory piece in the lifting from x_n to y_n of length t_n , and t_n goes to infinity. Since we have much larger control of pieces of trajectories of admissible type, we introduce also the *admissible rotation set* AR , where in the definition we consider only such pieces. Clearly, the admissible rotation set is contained in the rotation set. By the definition, both sets are closed. It is also clear that they are contained in the closed unit ball in \mathbb{R}^m , centered at the origin. Due to the time-reversibility, both sets R and AR are centrally symmetric with respect to the origin.

For a given point p in the phase space let us consider the trajectory $t \mapsto T(t)$ in \mathbb{R}^m starting at p . We can ask whether the limit of $(T(t) - T(0))/t$, as t goes to infinity, exists. If it does, we will call it the *rotation vector* of p . Clearly, it is the same for every point in the phase space of the full trajectory of p , so we can speak of the *rotation vector of a trajectory*. In particular, every periodic orbit has a rotation vector, and it is equal to $(T(s) - T(0))/s$, where s is the period of the orbit.

Note that if we use the discrete time (the number of reflections) rather than continuous time, we would get all good properties of the admissible rotation set from the description of the admissible sequences via the graph G and the results of [11]. Since we are using continuous time, the situation is more complicated. Nevertheless, Lemma 2.5 allows us to get similar results. For a trajectory piece T we will denote by $|T|$ its length and by $d(T)$ its displacement.

Theorem 3.1. *The admissible rotation set of a billiard on a torus with a small obstacle is convex.*

Proof. Fix vectors $u, v \in AR$ and a number $t \in (0, 1)$. We want to show that the vector $tu + (1 - t)v$ belongs to AR . Fix $\varepsilon > 0$. By the definition, there are finite admissible sequences A, B and trajectory pieces T, S of type A, B respectively, such that

$$\left\| \frac{d(T)}{|T|} - u \right\| < \varepsilon \quad \text{and} \quad \left\| \frac{d(S)}{|S|} - v \right\| < \varepsilon. \tag{3.1}$$

Both A, B can be represented as finite paths in the graph G . By Lemma 2.10, there are admissible sequences C_1, C_2, C_3 represented in G as paths of length at most 3, via elements of U , such that the concatenations of the form

$$D = AC_1AC_1 \dots AC_1AC_2BC_3BC_3 \dots BC_3B$$

are admissible. There exists a trajectory piece Q of type D . We will estimate its displacement and length.

Assume that in D the block A appears p times and the block B appears $q - p$ times. Let d_A, d_B be the total displacements due to the blocks A, B respectively. We get

$$\|d_A - pd(T)\| \leq 2pc \quad \text{and} \quad \|d_B - (q - p)d(S)\| \leq 2(q - p)c.$$

The displacement due to each of the blocks C_1, C_2, C_3 is at most of norm $2 + 2c$, so the total displacement due to all those blocks is at most of norm $q(2 + 2c)$. If we replace all displacements by the trajectory lengths, we get the same estimates (we use here Lemma 2.5). Thus we get the following estimates:

$$\|d(Q) - \alpha\| \leq 4qc + 2q \quad \text{and} \quad ||Q| - \beta| \leq 4qc + 2q,$$

where

$$\alpha = p d(T) + (q - p) d(S) \quad \text{and} \quad \beta = p|T| + (q - p)|S|.$$

Therefore

$$\begin{aligned} \left\| \frac{d(Q)}{|Q|} - \frac{\alpha}{\beta} \right\| &\leq \left\| \frac{d(Q)}{|Q|} - \frac{\alpha}{|Q|} \right\| + \left\| \frac{\alpha}{|Q|} - \frac{\alpha}{\beta} \right\| \\ &\leq \frac{4qc + 2q}{|Q|} + \|\alpha\| \frac{4qc + 2q}{|Q|\beta} \\ &= (4c + 2) \frac{q}{|Q|} \left(1 + \frac{\|\alpha\|}{\beta} \right). \end{aligned} \quad (3.2)$$

Set $s = p|T|/\beta$. Then $1 - s = (q - p)|S|/\beta$, so

$$\frac{\alpha}{\beta} = s \frac{d(T)}{|T|} + (1 - s) \frac{d(S)}{|S|}. \quad (3.3)$$

By (3.1), we get

$$\frac{\|\alpha\|}{\beta} \leq \max(\|u\|, \|v\|) + \varepsilon.$$

Moreover,

$$\frac{|Q|}{q} \geq \frac{\beta}{q} - (4c + 2) \geq \min(|T|, |S|) - (4c + 2).$$

Therefore if $|T|$ and $|S|$ are sufficiently large (we may assume this), the right-hand side of (3.2) is less than ε . Together with (3.3), we get

$$\left\| \frac{d(Q)}{|Q|} - \left(s \frac{d(T)}{|T|} + (1 - s) \frac{d(S)}{|S|} \right) \right\| < \varepsilon.$$

By this inequality and (3.1), it remains to show that by the right choice of p, q we can approximate t by s with an arbitrary accuracy.

We can write $s = f(x)$, where $x = p/q$ and

$$f(x) = \frac{|T|x}{|T|x + |S|(1 - x)}.$$

The function f is continuous on $[0, 1]$, takes value 0 at 0 and value 1 at 1. Therefore the image of the set of rational numbers from $(0, 1)$ is dense in $[0, 1]$. This completes the proof. \square

Theorem 3.2. *For a billiard on a torus with a small obstacle, rotation vectors of periodic orbits of admissible type are dense in the admissible rotation set.*

Proof. Fix a vector $\mathbf{u} \in AR$ and $\varepsilon > 0$. We want to find a periodic orbit of admissible type whose rotation vector is in the ε -neighborhood of \mathbf{u} . By the definition, there is an admissible sequence A and a trajectory piece T of type A such that

$$\left\| \frac{\mathbf{d}(T)}{|T|} - \mathbf{u} \right\| < \frac{\varepsilon}{2}. \tag{3.4}$$

Moreover, we can assume that $|T|$ is as large as we need. As in the proof of Theorem 3.1, we treat A as a path in the graph G and find an admissible sequence C represented in G as paths of length at most 3, via elements of U , such that the periodic concatenation $D = ACACAC\dots$ is admissible. There exists a periodic orbit of type D . Let Q be its piece corresponding to the itinerary AC . We will estimate its displacement and length.

Similarly as in the proof of Theorem 3.1, we get

$$|\mathbf{d}(Q) - \mathbf{d}(T)| \leq 4c + 2 \quad \text{and} \quad ||Q| - |T|| \leq 4c + 2.$$

Therefore

$$\begin{aligned} \left\| \frac{\mathbf{d}(Q)}{|Q|} - \frac{\mathbf{d}(T)}{|T|} \right\| &\leq \left\| \frac{\mathbf{d}(Q)}{|Q|} - \frac{\mathbf{d}(T)}{|Q|} \right\| + \left\| \frac{\mathbf{d}(T)}{|Q|} - \frac{\mathbf{d}(T)}{|T|} \right\| \\ &\leq \frac{4c + 2}{|T| - (4c + 2)} + \frac{\|\mathbf{d}(T)\|}{|T|} \cdot \frac{4c + 2}{|T| - (4c + 2)}. \end{aligned}$$

If $|T|$ is sufficiently large then the right-hand side of this inequality is smaller than $\varepsilon/2$. Together with (3.4) we get

$$\left\| \frac{\mathbf{d}(Q)}{|Q|} - \mathbf{u} \right\| < \varepsilon.$$

This completes the proof. \square

We will refer to closed paths in G as *loops*.

Remark 3.3. It is clear that in the above theorem we can additionally require that the corresponding loop in the graph G passes through a given vertex.

To get more results, we need a generalization of a lemma from [8] to higher dimensions.

Lemma 3.4. *Assume that $\mathbf{0} \in \mathbb{R}^m$ lies in the interior of the convex hull of a set of $m + 1$ vectors $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_m$. For every $K > 0$ if L is large enough then the following property holds. If $\mathbf{x} \in \mathbb{R}^m$ and $\|\mathbf{x}\| \leq L$ then there exists $i \in \{0, 1, \dots, m\}$ and a positive integer n such that $\|\mathbf{x} + n\mathbf{v}_i\| \leq L - K$. Moreover, $\|\mathbf{x} + j\mathbf{v}_i\| \leq L$ for $j = 1, 2, \dots, n - 1$.*

Proof. Let us fix $K > 0$. We will consider only L such that $L > K$. Set $M = \max_i \|\mathbf{v}_i\|$.

For each $\mathbf{x} \in \mathbb{R}^m$ with $\|\mathbf{x}\| = 1$ let $f(\mathbf{x})$ be the minimum of $\|\mathbf{x} + t\mathbf{v}_i\|$ over $i = 0, 1, \dots, m$ and $t \geq 0$. By the assumption, $f(\mathbf{x}) < 1$. Clearly f is continuous, and therefore there is $\varepsilon > 0$ such that $f(\mathbf{x}) \leq 1 - \varepsilon$ for every \mathbf{x} . Thus, for every $\mathbf{y} \in \mathbb{R}^m$ there is $i = 0, 1, \dots, m$ and $s \geq 0$ such that $\|\mathbf{y} + s\mathbf{v}_i\| \leq (1 - \varepsilon)\|\mathbf{y}\|$. Let n be the smallest integer larger than s . Then $n > 0$, and if $L \geq (M + K)/\varepsilon$ and $\|\mathbf{y}\| \leq L$ then $\|\mathbf{y} + n\mathbf{v}_i\| \leq (1 - \varepsilon)L + M \leq L - K$.

The last statement of the lemma follows from the convexity of the balls in \mathbb{R}^m . \square

Now we can follow the methods of [8] and [11]. We assume that our billiard has a small obstacle. For a full trajectory T we will denote by $T(t)$ the point to which we get after time t .

Lemma 3.5. *If \mathbf{u} is a vector from the interior of AR , then there exists a full trajectory T of admissible type and a constant M such that*

$$\|T(t) - T(0) - t\mathbf{u}\| \leq M \tag{3.5}$$

for all $t \in \mathbb{R}$.

Proof. Let us think first of positive t 's. Since \mathbf{u} is in the interior of AR , one can choose $m + 1$ vectors $\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m \in AR$ such that \mathbf{u} is in the interior of the convex hull of those vectors. Moreover, by Theorem 3.2 and Remark 3.3 we may assume that \mathbf{w}_i are rotation vectors of periodic orbits P_i of admissible type, corresponding to loops A_i in G passing through a common vertex V . We can also consider those loops as finite paths, ending at V and starting at the next vertex in the loop. Set

$$\mathbf{v}_i = \mathbf{d}(P_i) - |P_i|\mathbf{u} = |P_i|\mathbf{w}_i - |P_i|\mathbf{u} = |P_i|(\mathbf{w}_i - \mathbf{u}).$$

Since \mathbf{u} is in the interior of the convex hull of the vectors \mathbf{w}_i , we get that $\mathbf{0}$ is in the interior of the convex hull of the vectors $\mathbf{w}_i - \mathbf{u}$, and therefore $\mathbf{0}$ is in the interior of the convex hull of the vectors \mathbf{v}_i .

We will construct our trajectory, or rather a corresponding path in the graph G , by induction, using Lemma 3.4. Then we get a corresponding trajectory of admissible type by Theorem 2.2. We start with the empty sequence, that corresponds to the trajectory piece consisting of one point. Then, when a path B_j in G (corresponding to a trajectory piece Q_j) is constructed, and it ends at V , we look at the vector $\mathbf{x} = \mathbf{d}(Q_j) - |Q_j|\mathbf{u}$ and choose \mathbf{v}_i and n according to Lemma 3.4. We append B_j by adding n repetitions of A_i (corresponding to a trajectory piece that we can call nP_i) and obtain B_{j+1} (corresponding to a trajectory piece Q_{j+1}). To do all this, we have to define K that is used in Lemma 3.4 and prove that if $\|\mathbf{x}\| \leq L$ then also $\|\mathbf{d}(Q_{j+1}) - |Q_{j+1}|\mathbf{u}\| \leq L$.

Let us analyze the situation. When we concatenate Q_j and $A_i \dots A_i$ (n times) to get Q_{j+1} , by Lemma 2.5 we have

$$\|(\mathbf{d}(Q_{j+1}) - |Q_{j+1}|\mathbf{u}) - (\mathbf{d}(Q_j) - |Q_j|\mathbf{u}) - (\mathbf{d}(nP_i) - |nP_i|\mathbf{u})\| \leq 4c(1 + \|\mathbf{u}\|).$$

Moreover,

$$\mathbf{d}(nP_i) - |nP_i|\mathbf{u} = n(\mathbf{d}(P_i) - |P_i|\mathbf{u}) = n\mathbf{v}_i.$$

Therefore in Lemma 3.4 we have to take $K = 4c(1 + \|\mathbf{u}\|)$ and then we can make the induction step. In such a way we obtain an infinite path B in G . By Theorem 2.2, there exists a billiard trajectory T of type B .

Note that we did not complete the proof yet, because we got (3.5) (with $M = L$) only for a sequence of times $t = |Q_j|$. We can do better using the last statement of Lemma 3.4. This shows that (3.5) with $M = L + K$ holds for a sequence of times t with the difference of two consecutive terms of this sequence not exceeding $s = \max(|P_0|, |P_1|, \dots, |P_n|) + 4c$. Every time t' can be written as $t + r$ with t being a term of the above sequence (so that (3.5) holds with $M = L + K$) and $r \in [0, s)$. Then

$$\|T(t') - T(0) - t'\mathbf{u}\| \leq L + K + \|T(t+r) - T(t)\| + r\|\mathbf{u}\|.$$

Thus, (3.5) holds for all times with $M = L + K + s + s\|\mathbf{u}\|$.

The same can be done for negative t 's, so we get a full (two-sided) path, and consequently a full trajectory. \square

Now we are ready to prove the next important theorem. Remember that our phase space is a factor of a compact connected subset of the unit tangent bundle over the torus.

Theorem 3.6. *For a billiard on a torus with a small obstacle, if \mathbf{u} is a vector from the interior of AR , then there exists a compact invariant subset Y of the phase space, such that every trajectory from Y has admissible type and rotation vector \mathbf{u} .*

Proof. Let Y be the closure of the trajectory T from Lemma 3.5, taken in the phase space. If S is a trajectory obtained from T by starting it at time s (that is, $S(t) = T(s+t)$) then by Lemma 3.5 we get $\|S(t) - S(0) - t\mathbf{u}\| \leq 2M$ for all t . By continuity of the flow, this property extends to every trajectory S from Y . This proves that every trajectory from Y has rotation vector \mathbf{u} .

Since a trajectory of admissible type has no tangencies to the obstacle (by Condition 4 of the definition of admissible sequences), each finite piece of a trajectory from Y has admissible type. Therefore every trajectory from Y has admissible type. \square

Remark 3.7. The set Y above can be chosen minimal, and therefore the trajectory from Lemma 3.5 can be chosen recurrent.

As a trivial corollary to Theorem 3.6 we get the following.

Corollary 3.8. *For a billiard on a torus with a small obstacle, if \mathbf{u} is a vector from the interior of AR , then there exists a trajectory of admissible type with rotation vector \mathbf{u} .*

We also get another corollary, which follows from the existence of an ergodic measure on Y .

Corollary 3.9. *For a billiard on a torus with a small obstacle, if \mathbf{u} is a vector from the interior of AR , then there exists an ergodic invariant probability measure in the phase space, for which the integral of the velocity is equal to \mathbf{u} and almost every trajectory is of admissible type.*

This corollary is stronger than Corollary 3.8, because from it and from the Ergodic Theorem it follows that almost every point has rotation vector \mathbf{u} . The details of the necessary formalism are described in Sect. 6. Of course, in our particular case both results are corollaries to Theorem 3.6, so we know anyway that all points of Y have rotation vector \mathbf{u} .

4. Admissible Rotation Set is Large

In this section we will investigate how large the admissible rotation set AR is. This of course depends on the size of the obstacle and the dimension of the space. We will measure the size of AR by the radius of the largest ball centered at the origin and contained in AR .

We will start with the estimates that depend on the dimension m of the space but not on the size of the obstacle (provided it is small in our meaning). In order to do it, we first identify some elements of \mathbb{Z}^m that are always vertices of G . Recall that

$$A_m = \{-1, 0, 1\}^m \setminus \{\mathbf{0}\}.$$

Lemma 4.1. *If $\mathbf{k} \in A_m$ then $(\sqrt{2}/2)(\mathbf{k}/\|\mathbf{k}\|) \in AR$.*

Proof. If $\mathbf{k} \in U$ then there is a vector $\mathbf{l} \in U$ orthogonal to \mathbf{k} . Vectors $\mathbf{k} + \mathbf{l}$ and $\mathbf{k} - \mathbf{l}$ belong to A_m and one can easily check that there are edges from $\mathbf{k} + \mathbf{l}$ to $\mathbf{k} - \mathbf{l}$ and from $\mathbf{k} - \mathbf{l}$ to $\mathbf{k} + \mathbf{l}$ in G . The periodic path $(\mathbf{k} + \mathbf{l})(\mathbf{k} - \mathbf{l})(\mathbf{k} + \mathbf{l})(\mathbf{k} - \mathbf{l}) \dots$ in G gives us a periodic orbit P of the billiard. The displacement along P is $2\mathbf{k}$ and the period of P is smaller than $\|\mathbf{k} + \mathbf{l}\| + \|\mathbf{k} - \mathbf{l}\| = 2\sqrt{2}$, so the rotation vector of P is $t\mathbf{k}$, where $t > \sqrt{2}/2$. Since $\mathbf{0} \in AR$ and AR is convex, we get $(\sqrt{2}/2)(\mathbf{k}/\|\mathbf{k}\|) \in AR$.

Assume now that $\mathbf{k} \in A_m$ and $\|\mathbf{k}\| > 1$. Then $\mathbf{k} = \mathbf{l} + \mathbf{u}$ for some $\mathbf{l} \in A_m$ and $\mathbf{u} \in U$ such that \mathbf{u} is orthogonal to \mathbf{l} . By Lemma 2.9, \mathbf{l} is a vertex of G . By Lemma 2.8 there are edges in G from \mathbf{l} to \mathbf{u} and from \mathbf{u} to \mathbf{l} . Similarly as before, we get a periodic orbit of the billiard (corresponding to the periodic path $\mathbf{l}\mathbf{u}\mathbf{l}\mathbf{u} \dots$) with the displacement \mathbf{k} and period less than $\|\mathbf{l}\| + \|\mathbf{u}\| = \sqrt{\|\mathbf{k}\|^2 - 1} + 1$, so

$$\frac{\|\mathbf{k}\|}{\sqrt{\|\mathbf{k}\|^2 - 1} + 1} \cdot \frac{\mathbf{k}}{\|\mathbf{k}\|} \in AR.$$

Since $\|\mathbf{k}\|/(\sqrt{\|\mathbf{k}\|^2 - 1} + 1) \geq \sqrt{2}/2$, the vector $(\sqrt{2}/2)(\mathbf{k}/\|\mathbf{k}\|)$ also belongs to AR . \square

By the results of [2], the convex hull of $\{\mathbf{v}/\|\mathbf{v}\| : \mathbf{v} \in A_m\}$ contains the closed ball centered at $\mathbf{0}$ with radius $2/\sqrt{\ln m + 5}$. From this and Lemma 4.1 we get immediately the following result.

Theorem 4.2. *For a billiard on a torus with a small obstacle, the set AR contains the closed ball centered at $\mathbf{0}$ with radius $\sqrt{2}/(\ln m + 5)$.*

Now we proceed to the estimates that are independent of the dimension m . This is not as simple as it seems. As we saw above, a straightforward attempt that takes into account only those vectors of \mathbb{Z}^m for which we can show explicitly that they are vertices of G , gives estimates that go to 0 as $m \rightarrow \infty$. By the results of [2], those estimates cannot be significantly improved. Therefore we have to use another method.

Let us assume first that O_0 is the ball centered at $\mathbf{0}$ of radius $r < \sqrt{2}/4$. We start with a simple lemma.

Lemma 4.3. *Assume that O_l is between O_0 and O_k and let ϑ be the angle between the vectors \mathbf{k} and \mathbf{l} . Then*

$$\langle \mathbf{k}, \mathbf{l} \rangle^2 \geq \|\mathbf{k}\|^2 \left(\|\mathbf{l}\|^2 - 4r^2 \right) > \|\mathbf{k}\|^2 \left(\|\mathbf{l}\|^2 - \frac{1}{2} \right) \tag{4.1}$$

and

$$\sin \vartheta \leq 2r/\|\mathbf{l}\|. \tag{4.2}$$

Proof. If O_l is between O_0 and O_k then there is a line parallel to the vector \mathbf{k} , whose distances from $\mathbf{0}$ and \mathbf{l} are at most r . Therefore the distance of \mathbf{l} from the line through $\mathbf{0}$ and \mathbf{k} is at most $2r$. The orthogonal projection of \mathbf{l} to this line is $(\langle \mathbf{k}, \mathbf{l} \rangle / \|\mathbf{k}\|^2)\mathbf{k}$, so

$$\left\| \mathbf{l} - \frac{\langle \mathbf{k}, \mathbf{l} \rangle}{\|\mathbf{k}\|^2} \mathbf{k} \right\|^2 \leq 4r^2.$$

The left-hand side of this inequality is equal to

$$\|I\|^2 - \frac{\langle k, I \rangle^2}{\|k\|^2}$$

and $4r^2 < 1/2$, so (4.1) holds.

By (4.1), we have

$$\sin^2 \vartheta = 1 - \cos^2 \vartheta = 1 - \frac{\langle k, I \rangle^2}{\|k\|^2 \|I\|^2} \leq 1 - \frac{\|k\|^2 (\|I\|^2 - 4r^2)}{\|k\|^2 \|I\|^2} = \frac{4r^2}{\|I\|^2},$$

so (4.2) holds. \square

Clearly, the angle ϑ above is acute.

The estimate in the next lemma requires extensive use of the fact that the vectors that we are considering have integer components.

Lemma 4.4. *Assume that O_I is between O_0 and O_k and*

$$\langle k, I \rangle \leq \langle k, k - I \rangle.$$

Then $\|I\| \leq \|k\|/2$.

Proof. By Lemma 4.3, (4.1) holds. Since $\langle k, I \rangle \leq \langle k, k - I \rangle$, we get $\langle k, I \rangle \leq \|k\|^2 - \langle k, I \rangle$, so

$$2\langle k, I \rangle \leq \|k\|^2. \tag{4.3}$$

By (4.1) and (4.3) we get

$$\|k\|^2 > 4\|I\|^2 - 2. \tag{4.4}$$

If $\|I\| > \|k\|/2$, then $\|k\|^2 < 4\|I\|^2$. Together with (4.4), since $\|k\|^2$ and $\|I\|^2$ are integers, we get

$$\|k\|^2 = 4\|I\|^2 - 1. \tag{4.5}$$

From (4.3) and (4.5) we get $\langle k, I \rangle \leq 2\|I\|^2 - 1/2$. Hence, since $\langle k, I \rangle$ is also an integer, we get $\langle k, I \rangle \leq 2\|I\|^2 - 1$. From this, (4.1) and (4.5) we get

$$(2\|I\|^2 - 1)^2 > (4\|I\|^2 - 1) \left(\|I\|^2 - \frac{1}{2} \right) = \left(2\|I\|^2 - \frac{1}{2} \right) (2\|I\|^2 - 1),$$

a contradiction. \square

Let us think about standing at the origin and looking at the sky, where vertices of G are stars. Are there big parts of the sky without a single star? Observe that as the dimension m of the space grows, the angles between the integer vectors tend to become larger. For instance, the angle between the vectors $(1, 0, \dots, 0)$ and $(1, \dots, 1)$ is of order $\pi/2 - 1/\sqrt{m}$. Thus, any given acute angle can be considered relatively small if m is sufficiently large.

Let α be a positive angle. We will say that a set $A \subset \mathbb{Z}^m \setminus \{\mathbf{0}\}$ is α -dense in the sky if for every $v \in \mathbb{R}^m \setminus \{\mathbf{0}\}$ there is $u \in A$ such that the angle between the vectors v and u is at most α . Set

$$\eta(r) = \sum_{n=0}^{\infty} \arcsin \frac{r}{2^{n-1}}.$$

Proposition 4.5. *The set of vertices of G is $\eta(r)$ -dense in the sky.*

Proof. Fix a vector $v \in \mathbb{R}^m \setminus \{0\}$ and $\varepsilon > 0$. There exists $k_0 \in \mathbb{Z}^m \setminus \{0\}$ such that the angle between v and k_0 is less than ε . Then we define by induction a finite sequence (k_1, k_2, \dots, k_n) of elements of $\mathbb{Z}^m \setminus \{0\}$ such that $O_{k_{i+1}}$ is between O_0 and O_{k_i} and $\|k_{i+1}\| \leq \|k_i\|/2$. This is possible by Lemmas 2.7 and 4.4. Since $\|k_i\| \geq 1$ for each i , this procedure has to terminate at some k_n . Then there is no obstacle between O_0 and O_{k_n} , so k_n is a vertex of G . By Lemma 4.3, the angle between k_i and k_{i+1} is at most $\arcsin(2r/\|k_{i+1}\|)$. By our construction, we have $\|k_n\| \geq 1 = 2^0$, $\|k_{n-1}\| \geq 2^1$, $\|k_{n-2}\| \geq 2^2$, etc. Therefore the angle between k_n and k_0 is smaller than $\eta(r)$. Hence, the angle between v and k_n is smaller than $\eta(r) + \varepsilon$. Since ε was arbitrary, this angle is at most $\eta(r)$. \square

Remark 4.6. Proposition 4.5 was proved under the assumption that O_0 is the ball centered at 0 of radius $r < \sqrt{2}/4$. However, making an obstacle smaller results in preservation or even enlargement of G . Moreover, we have a freedom in the lifting where to put the origin. Therefore, Proposition 4.5 remains true under a weaker assumption, that O is contained in a closed ball of radius $r < \sqrt{2}/4$.

Let us investigate the properties of $\eta(r)$.

Lemma 4.7. *The function η is continuous and increasing on $(0, \sqrt{2}/4]$. Moreover, $\eta(r) < \sqrt{2} \pi r$. In particular, $\eta(\sqrt{2}/4) < \pi/2$ and*

$$\lim_{r \rightarrow 0} \eta(r) = 0.$$

Proof. Assume that $0 < r \leq \sqrt{2}/4$. Then all numbers whose arcus sine we are taking are from the interval $(0, \sqrt{2}/2]$, so clearly η is continuous and increasing. We have also the estimate

$$\frac{x}{\arcsin x} \geq \frac{\sqrt{2}/2}{\pi/4} = \frac{2\sqrt{2}}{\pi}.$$

Moreover, the equality holds only if $x = \sqrt{2}/2$. Thus

$$\eta(r) < \sum_{n=0}^{\infty} \frac{\pi r}{2^n \sqrt{2}} = \sqrt{2} \pi r.$$

Therefore $\lim_{r \rightarrow 0} \eta(r) = 0$ and

$$\eta(\sqrt{2}/4) < \sqrt{2} \pi \frac{\sqrt{2}}{4} = \frac{\pi}{2}.$$

\square

Now we assume only that O is small. In the next lemma we obtain two estimates of the length of $t\mathbf{k} \in AR$ if \mathbf{k} is a vertex of G . One of those estimates will be useful for all vertices of G , the other one for those with large norm. The main idea of the proof is similar as in the proof of Lemma 4.1.

Lemma 4.8. *If \mathbf{k} is a vertex of G then the vectors $(1 - \sqrt{2}/2)(\mathbf{k}/\|\mathbf{k}\|)$ and $((\|\mathbf{k}\| - 1)/(\|\mathbf{k}\| + 1))(\mathbf{k}/\|\mathbf{k}\|)$ belong to AR .*

Proof. Let $\mathbf{k} = (x_1, x_2, \dots, x_m)$ be a vertex of G . Let s be the number of non-zero components of \mathbf{k} . Then we may assume that $x_i \neq 0$ if $i \leq s$ and $x_i = 0$ if $i > s$. If $s = 1$ then the statement of the lemma follows from Lemma 4.1.

Assume now that $s > 1$. Then for every $i \leq s$ there is a vector $\mathbf{v}_i \in U$ with only i^{th} component non-zero and $\langle \mathbf{v}_i, \mathbf{k} \rangle < 0$. By Lemma 2.8 there are edges in G from \mathbf{k} to \mathbf{v}_i and from \mathbf{v}_i to \mathbf{k} , so the periodic path $\mathbf{k}\mathbf{v}_i\mathbf{k}\mathbf{v}_i \dots$ in G gives us a periodic orbit P_i of the billiard. The displacement along P_i is $\mathbf{k} + \mathbf{v}_i$ and the period of P_i is smaller than $\|\mathbf{k}\| + \|\mathbf{v}_i\| = \|\mathbf{k}\| + 1$, so the rotation vector of P_i is $t_i(\mathbf{k} + \mathbf{v}_i)$ with $t_i > 1/(\|\mathbf{k}\| + 1)$. Therefore the vector $(\mathbf{k} + \mathbf{v}_i)/(\|\mathbf{k}\| + 1)$ belongs to AR .

Since the vectors \mathbf{v}_i form an orthonormal basis of \mathbb{R}^s , we have

$$\mathbf{k} = \sum_{i=1}^s \langle \mathbf{v}_i, \mathbf{k} \rangle \mathbf{v}_i.$$

Set

$$a = \sum_{i=1}^s \langle \mathbf{v}_i, \mathbf{k} \rangle \text{ and } a_i = \frac{\langle \mathbf{v}_i, \mathbf{k} \rangle}{a}.$$

Then the vector

$$\mathbf{u} = \sum_{i=1}^s a_i \frac{\mathbf{k} + \mathbf{v}_i}{\|\mathbf{k}\| + 1}$$

is a convex combination of elements of AR , so $\mathbf{u} \in AR$. We have

$$\mathbf{u} = \frac{\mathbf{k}}{\|\mathbf{k}\| + 1} \sum_{i=1}^s a_i + \frac{1}{a(\|\mathbf{k}\| + 1)} \sum_{i=1}^s \langle \mathbf{v}_i, \mathbf{k} \rangle \mathbf{v}_i = \frac{\mathbf{k}}{\|\mathbf{k}\| + 1} \left(1 + \frac{1}{a} \right).$$

For each i we have $\langle \mathbf{v}_i, \mathbf{k} \rangle \leq -1$, so $a \leq -s$, and therefore $1 + 1/a \geq (s - 1)/s$. Moreover,

$$\frac{\|\mathbf{k}\|}{\|\mathbf{k}\| + 1} \geq \frac{\sqrt{s}}{\sqrt{s} + 1}.$$

Since $s \geq 2$, we get

$$\frac{s - 1}{s} \cdot \frac{\sqrt{s}}{\sqrt{s} + 1} = \frac{\sqrt{s} - 1}{\sqrt{s}} \geq \frac{\sqrt{2} - 1}{\sqrt{2}} = 1 - \frac{\sqrt{2}}{2},$$

so the vector \mathbf{u} has the direction of \mathbf{k} and length at least $1 - \sqrt{2}/2$.

To get the other estimate of the length of \mathbf{u} , note that $\langle \mathbf{v}_i, \mathbf{k} \rangle = -|x_i|$, so

$$a = - \sum_{i=1}^s |x_i| \leq -\|\mathbf{k}\|,$$

and hence

$$\|\mathbf{u}\| \geq \frac{\|\mathbf{k}\|}{\|\mathbf{k}\| + 1} \cdot \left(1 - \frac{1}{\|\mathbf{k}\|} \right) = \frac{\|\mathbf{k}\| - 1}{\|\mathbf{k}\| + 1}.$$

□

Lemma 4.9. *Let $A \subset \mathbb{R}^m$ be a finite set, α -dense in the sky for some $\alpha < \pi/2$. Assume that every vector of A has norm c . Then the convex hull of A contains a ball of radius $c \cos \alpha$, centered at $\mathbf{0}$.*

Proof. Let K be the convex hull of A . Then K is a convex polytope with vertices from A . Since A is α -dense in the sky and $\alpha < \pi/2$, K is non-degenerate and $\mathbf{0}$ belongs to its interior. Let s be the radius of the largest ball centered at $\mathbf{0}$ and contained in K . This ball is tangent to some face of K at a point \mathbf{v} . Then the whole K is contained in the half-space $\{\mathbf{u} : \langle \mathbf{u}, \mathbf{v} \rangle \leq \|\mathbf{v}\|^2\}$. In particular, for every $\mathbf{u} \in A$ we have $\langle \mathbf{u}, \mathbf{v} \rangle \leq \|\mathbf{v}\|^2$. Since A is α -dense in the sky, there is $\mathbf{u} \in A$ such that the angle between \mathbf{v} and \mathbf{u} is at most α . Therefore

$$\|\mathbf{v}\|^2 \geq \langle \mathbf{u}, \mathbf{v} \rangle \geq \|\mathbf{u}\| \|\mathbf{v}\| \cos \alpha = c \|\mathbf{v}\| \cos \alpha,$$

so $s = \|\mathbf{v}\| \geq c \cos \alpha$. \square

Now we can get the first, explicit, estimate of the radius of the largest ball contained in AR .

Theorem 4.10. *For a billiard on a torus with small obstacle, assume that O is contained in a closed ball of radius $r < \sqrt{2}/4$. Then the admissible rotation set contains the closed ball of radius $(1 - \sqrt{2}/2) \cos \eta(r)$ centered at $\mathbf{0}$.*

Proof. Set $H = \{(1 - \sqrt{2}/2)(\mathbf{k}/\|\mathbf{k}\|) : \mathbf{k} \text{ is a vertex of } G\}$ and let K be the convex hull of H . By Lemma 4.8 and by the convexity of AR , we have $K \subset AR$. By Proposition 4.5, H is $\eta(r)$ -dense in the sky. Thus, by Lemma 4.9, K contains the closed ball of radius $(1 - \sqrt{2}/2) \cos \eta(r)$ centered at $\mathbf{0}$. \square

The second estimate is better for small r (uniformly in m), but does not give an explicit formula for the radius of the ball contained in AR . We first need two lemmas.

Lemma 4.11. *For every integer $N > 1$ there exists an angle $\beta(N) > 0$ (independent of m), such that if $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^m \setminus \mathbf{0}$ are vectors of norm less than N and the angle ϑ between \mathbf{u} and \mathbf{v} is positive, then $\vartheta \geq \beta(N)$.*

Proof. Under our assumptions, each of the vectors \mathbf{u}, \mathbf{v} has less than N^2 non-zero components. Therefore the angle between \mathbf{u} and \mathbf{v} is the same as the angle between some vectors $\mathbf{u}', \mathbf{v}' \in \mathbb{Z}^{2N^2-2}$ of the same norms as \mathbf{u}, \mathbf{v} . However, there are only finitely many vectors in $\mathbb{Z}^{2N^2-2} \setminus \{\mathbf{0}\}$ of norm less than N , so the lemma holds with $\beta(N)$ equal to the smallest positive angle between such vectors. \square

Lemma 4.12. *Let $A \subset \mathbb{Z}^m \setminus \mathbf{0}$ be a finite set, α -dense in the sky for some $\alpha < \beta(N)/2$, where $\beta(N)$ is as in Lemma 4.11. Then the set of those elements of A which have norm at least N is 2α -dense in the sky.*

Proof. Let B be the set of vectors of $\mathbb{R}^m \setminus \{\mathbf{0}\}$ whose angular distance from some vector of A of norm at least N is 2α or less.

Let $\mathbf{v} \in \mathbb{R}^m$ be a non-zero vector. Assume that the angles between \mathbf{v} and all vectors of A are non-zero. By the assumptions, there exists $\mathbf{u} \in A$ such that the angle ϑ between \mathbf{v} and \mathbf{u} is at most α . If $\|\mathbf{u}\| \geq N$ then $\mathbf{v} \in B$. Suppose that $\|\mathbf{u}\| < N$. Choose $\varepsilon > 0$ such that $\varepsilon < \beta(N) - 2\alpha$ and $\varepsilon < \vartheta$. We draw a great circle in the sky through \mathbf{u} and \mathbf{v} and go along it from \mathbf{u} through \mathbf{v} and beyond it to some \mathbf{v}' so that the angle between

\mathbf{u} and \mathbf{v}' is $\alpha + \varepsilon$. Now, there exists $\mathbf{u}' \in A$ such that the angle between \mathbf{v}' and \mathbf{u}' is at most α . Then the angle between \mathbf{u} and \mathbf{u}' is at least ε and at most $2\alpha + \varepsilon < \beta(N)$, so $\|\mathbf{u}'\| \geq N$. The angle between \mathbf{v} and \mathbf{u}' is at most $2\alpha + \varepsilon - \vartheta < 2\alpha$, so again, $\mathbf{v} \in B$.

In such a way we have shown that B is dense in $\mathbb{R}^m \setminus \{\mathbf{0}\}$. It is also clearly closed in $\mathbb{R}^m \setminus \{\mathbf{0}\}$, so it is equal to $\mathbb{R}^m \setminus \{\mathbf{0}\}$. \square

Let $\beta(N)$ be as in Lemma 4.11, and let $N(r)$ be the maximal N such that $\eta(r) < \beta(N)/2$. This definition is correct for sufficiently small r , since clearly $\beta(N) \rightarrow 0$ as $N \rightarrow \infty$, and by Lemma 4.7, $\eta(r) \rightarrow 0$ as $r \rightarrow 0$. It follows that $N(r) \rightarrow \infty$ as $r \rightarrow 0$.

Theorem 4.13. *For a billiard on a torus with a small obstacle, assume that O is contained in a closed ball of radius $r < \sqrt{2}/4$ for r so small that $N(r)$ is defined. Then the admissible rotation set contains the closed ball of radius $((N(r) - 1)/(N(r) + 1)) \cos(2\eta(r))$ centered at $\mathbf{0}$.*

Proof. Since $\eta(r) < \beta(N(r))/2$, by Proposition 4.5, Remark 4.6 and Lemma 4.12, the set of those vertices of G that have norm at least $N(r)$ is $2\eta(r)$ -dense in the sky. By Lemma 4.8, if \mathbf{k} is such a vertex, $((\|\mathbf{k}\| - 1)/(\|\mathbf{k}\| + 1))(\mathbf{k}/\|\mathbf{k}\|) \in AR$. Since $\mathbf{0} \in AR$ and AR is convex, also $((N(r) - 1)/(N(r) + 1))(\mathbf{k}/\|\mathbf{k}\|) \in AR$. Then by Lemma 4.9, the closed ball of radius $((N(r) - 1)/(N(r) + 1)) \cos(2\eta(r))$ centered at $\mathbf{0}$ is contained in AR . \square

Corollary 4.14. *The radius of the largest ball centered at $\mathbf{0}$ contained in the admissible rotation set goes to 1 uniformly in m as the diameter of the obstacle goes to 0.*

We conclude this section with a result showing that even though AR may be large, it is still smaller than R .

Theorem 4.15. *For a billiard on a torus with a small obstacle, the admissible rotation set is contained in the open unit ball. In particular, $AR \neq R$.*

Proof. Since the graph G is finite, there exist positive constants $c_1 < c_2$ such that for every trajectory piece of admissible type the distance between two consecutive reflections is contained in $[c_1, c_2]$. Moreover, from the definition of an edge in G and from the compactness of the obstacle it follows that there is an angle $\alpha > 0$ such that the direction of a trajectory piece of admissible type changes by at least α at each reflection. Consider the triangle with two sides of length c_1 and c_2 and the angle $\pi - \alpha$ between them. Let a be the ratio between the length of the third side and $c_1 + c_2$, that is,

$$a = \frac{\sqrt{c_1^2 + c_2^2 + 2c_1c_2 \cos \alpha}}{c_1 + c_2}.$$

This ratio is less than 1 and it decreases when α or c_1/c_2 grows. Therefore, if consecutive reflections for a trajectory piece of admissible type are at times t_1, t_2, t_3 , then the displacement between the first and the third reflections divided by $t_3 - t_1$ is at most a . Thus, every vector from AR has length at most a .

Clearly, the vector $(1, 0, \dots, 0)$ belongs to R , and thus $AR \neq R$. \square

5. Billiard in the Square

Now we consider a billiard in the square $S = [-1/2, 1/2]^2$ with one convex obstacle O with a smooth boundary. The lifting to \mathbb{R}^2 , considered in Sect. 2 is replaced in this case by the unfolding to \mathbb{R}^2 . That is, we cover \mathbb{R}^2 by the copies of S obtained by consecutive symmetries with respect to the lines $x = n + 1/2$ and $y = n + 1/2$, $n \in \mathbb{Z}$. Thus, the square $S_{\mathbf{k}} = S + \mathbf{k}$ ($\mathbf{k} \in \mathbb{Z}^2$) with the obstacle $O_{\mathbf{k}}$ in it is the square S with O , translated by \mathbf{k} , with perhaps an additional symmetry applied. If $\mathbf{k} = (p, q)$, then, if both p, q are even, there is no additional symmetry; if p is even and q odd, we apply symmetry with respect to the line $y = q$; if p is odd and q is even, we apply symmetry with respect to the line $x = p$; and if both p, q are odd, we apply central symmetry with respect to the point (p, q) . In this model, trajectories in S with obstacle O unfold to trajectories in \mathbb{R}^2 with obstacles $O_{\mathbf{k}}$, $\mathbf{k} \in \mathbb{Z}^2$.

The situation in \mathbb{R}^2 is now the same as in the case of the torus billiard, except that, as we mentioned above, the obstacles are not necessarily the translations of O_0 , and, of course, the observable whose averages we take to get the rotation set is completely different. Let us trace which definitions, results and proofs of Sects. 2, 3 and 4 remain the same, and which need modifications.

The definitions of *between* and *type* remain the same. Lemma 2.1 is still valid, but in its proof we have to look at the trajectory on the torus $\mathbb{R}^2/(2\mathbb{Z})^2$ rather than $\mathbb{R}^2/\mathbb{Z}^2$. Then the definition of an *admissible sequence* remains the same.

The first part of Theorem 2.2 and its proof remains the same, but in the proof of the part about periodic trajectories we have to be careful. The point $\mathbf{x}_0 + \mathbf{p}$ from the last paragraph of the proof has to be replaced by a point that after folding (the operation reverse to the unfolding) becomes \mathbf{x}_0 . This gives us a periodic orbit in the unfolding that projects (folds) to a periodic orbit in the square. Moreover, there may be a slight difference between the square case and the torus case if we want to determine the least discrete period of this orbit (where in the square case, in analogy to the torus case, we count only reflections from the obstacle). In the torus case it is clearly the same as the least period of the type. In the square case this is not necessarily so. For instance, if the obstacle is a disk centered at the origin, the orbit that goes vertically from the highest point of the disk, reflects from the upper side of the square and returns to the highest point of the disk, has discrete period 1 in the above sense. However, its type is periodic of period 2 and in the unfolding it has period 2. Fortunately, such things are irrelevant for the rest of our results.

Lemma 2.3, Corollary 2.4 and their proofs remain the same as in the torus case. The same can be said about the part of Lemma 2.5 that refers to the lengths of trajectory pieces.

The definition of the graph G has to be modified. This is due to the fact that the conditions (3) and (4) cannot be restated in the same way as in the torus case, because now not only translations, but also symmetries are involved (the obstacle needs not be symmetric, and the unfolding process involves symmetries about vertical and horizontal lines). In order to eliminate symmetries, we enlarge the number of vertices of G four times. Instead of O_0 , we look at $\{O_0, O_{(1,0)}, O_{(0,1)}, O_{(1,1)}\}$. For every $\mathbf{k} = (p, q) \in \mathbb{Z}^2$ there is $\zeta(\mathbf{k}) \in Q$, where $Q = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$, such that $\mathbf{k} - \zeta(\mathbf{k})$ has both components even. Then condition (3) can be restated as no obstacle between $O_{\zeta(\mathbf{k}_{n-1})}$ and $O_{I_n + \zeta(\mathbf{k}_{n-1})}$, and condition (4) as the obstacle $O_{I_n + \zeta(\mathbf{k}_{n-1})}$ not between $O_{\zeta(\mathbf{k}_{n-1})}$ and $O_{I_n + I_{n+1} + \zeta(\mathbf{k}_{n-1})}$. Therefore we can take as the vertices of G the pairs (i, j) , where $\mathbf{i} \in Q$, $\mathbf{j} \in \mathbb{Z}^2$, $\mathbf{i} \neq \mathbf{j}$, and there is no obstacle between $O_{\mathbf{i}}$ and $O_{\mathbf{j}}$. There is an edge in G from (\mathbf{i}, \mathbf{j}) to $(\mathbf{i}', \mathbf{j}')$ if and only if $O_{\mathbf{j}}$ is not between $O_{\mathbf{i}}$ and $O_{\mathbf{j} + \mathbf{j}' - \mathbf{i}'}$ and $\zeta(\mathbf{j}) = \mathbf{i}'$. Then,

similarly as in the torus case, there is a one-to-one correspondence between admissible sequences and one-sided infinite paths in G , starting at vertices $(\mathbf{0}, j)$.

This restriction on the starting point of a path in G creates some complication, but for every $i \in Q$ there is a similar correspondence between admissible sequences and one-sided infinite paths in G , starting at vertices (i, j) . Therefore, if we want to glue finite paths, we may choose an appropriate i .

Lemma 2.6 and its proof remain unchanged. The definition of a *small obstacle* has to be modified slightly. This is due to the fact, that while a torus is homogeneous, so all positions of an obstacle are equivalent, this is not the case for a square. An obstacle placed close to a side of the square will produce a pattern of obstacles in the unfolding which is difficult to control. Therefore we will say that the obstacle O is *small* if it is contained in a closed ball of radius smaller than $\sqrt{2}/4$, centered at $\mathbf{0}$. With this definition, Lemmas 2.8 and 2.10 and their proofs remain unchanged.

We arrived at a point where the situation is completely different than for the torus case, namely, we have to define the displacement function. Once we do it, the definitions of the rotation set, admissible rotation set and rotation vector remain the same, except that we will call a rotation vector (since it belongs to \mathbb{R}) a *rotation number*.

Since we have to count how many times the trajectory rotates around the obstacle, the simplest way is to choose a point z in the interior of O and set $\varphi(x) = \arg(x - z)/(2\pi)$, where \arg is the complex argument (here we identify \mathbb{R}^2 with \mathbb{C}). It is not important that the argument is multivalued, since we are interested only in its increment along curves. For any closed curve Γ avoiding the interior of O , the increment of φ is equal to the winding number of Γ with respect to z . Since the whole interior of O lies in the same component of $\mathbb{C} \setminus \Gamma$, this number does not depend on the choice of z . If Γ is not closed, we can extend it to a closed one, while changing the increment of φ by less than 1. Therefore changing z will amount to the change of the increment of φ by less than 2. When computing the rotation numbers, we divide the increment of φ by the length of the trajectory piece, and this length goes to infinity. Therefore in the limit a different choice of z will give the same result. This proves that the rotation set we get is independent of the choice of z .

The proofs of the results of Sect. 3 rely on the second part of Lemma 2.5, which we have not discussed yet. The possibility of the trajectory pieces crossing the obstacles, mentioned in that lemma, was necessary only for the proof of its first part, so we do not have to worry about it now. However, we have to make an additional assumption that B is admissible. This creates no problem either, because this is how we use it later. Since we changed here many details, it makes sense to state exactly what we will be proving.

Lemma 5.1. *For every finite admissible sequence $B = (k_n)_{n=0}^s$ of elements of \mathbb{Z}^2 the displacements φ along trajectory pieces of type B differ by less than 2.*

Proof. Note that the “folding” map $\pi: \overline{\mathbb{R}^2 \setminus \bigcup_{k \in \mathbb{Z}^2} O_k} \rightarrow \overline{S \setminus O}$ is continuous. Therefore if curves Γ and γ in $\overline{\mathbb{R}^2 \setminus \bigcup_{k \in \mathbb{Z}^2} O_k}$ with the common beginning and common end are homotopic then $\pi(\Gamma)$ and $\pi(\gamma)$ are homotopic, so the increments of φ along them are the same. If Γ and Γ' are two trajectory pieces as in the statement of the lemma, then Γ' can be extended to γ with the same beginning and end as Γ , with the change of the increment of φ along its projection by π less than 2 (1 at the beginning, 1 at the end). Thus, it suffices to show that this can be done in such a way that Γ and γ are homotopic.

Therefore we have to analyze what may be the reasons for Γ and γ not to be homotopic. Extending Γ' to γ can be done in the right way, so this leaves two possibly bad things that we have to exclude. The first is that when going from O_{k_i} to $O_{k_{i+2}}$ via $O_{k_{i+1}}$,

we pass with Γ on one side of $O_{k_{i+1}}$ and with Γ' on the other side of $O_{k_{i+1}}$. The second one is that when going from O_{k_i} to $O_{k_{i+1}}$, we pass with Γ on one side of some O_j , and with Γ' on the other side of O_j . However, the first possibility contradicts condition (4) from the definition of admissibility and the second one contradicts condition (3). This completes the proof. \square

With Lemma 5.1 replacing Lemma 2.5, the rest of results of Sect. 3 (except the last two theorems and the corollary) and their proofs remain the same (with obvious minor modifications, for instance due to the fact that constants in Lemmas 2.5 and 5.1 are different). Let us state the main theorems we get in this way.

Theorem 5.2. *The admissible rotation set of a billiard in a square with a small obstacle is convex, and consequently, it is a closed interval symmetric with respect to 0.*

Theorem 5.3. *For a billiard in a square with a small obstacle, rotation numbers of periodic orbits of admissible type are dense in the admissible rotation set.*

Theorem 5.4. *For a billiard in a square with a small obstacle, if u is a number from the interior of AR , then there exists a compact, forward invariant subset Y of the phase space, such that every trajectory from Y has admissible type and rotation number u .*

Corollary 5.5. *For a billiard in a square with a small obstacle, if u is a number from the interior of AR , then there exists a trajectory of admissible type with rotation number u .*

Corollary 5.6. *For a billiard in a square with a small obstacle, if u is a number from the interior of AR , then there exists an ergodic invariant probability measure in the phase space, for which the integral of the displacement is equal to u and almost every trajectory is of admissible type.*

Since the billiard in the square is defined only in dimension 2, most of the results of Sect. 4 do not have counterparts here. However, we can investigate what happens to AR as the size of the obstacle decreases. Moreover, here the position of the obstacle matters, so the size of the obstacle should be measured by the radius of the smallest ball centered in the origin that contains it. The following theorem should be considered in the context of Theorem 6.9 that states that the full rotation set, R , is equal to $[-\sqrt{2}/4, \sqrt{2}/4]$.

Theorem 5.7. *For every $\varepsilon > 0$ there exists $\delta > 0$ such that the set AR contains the interval $[-\sqrt{2}/4 + \varepsilon, \sqrt{2}/4 - \varepsilon]$ whenever the obstacle is contained in the disk centered at the origin and diameter less than δ .*

Proof. Let us estimate the rotation number of the curve V that in the unfolding is a straight line segment from the origin to the point $(2n + 1, 2n)$. We are counting the displacement as the rotation around the center of the square (as always, as the multiples of 2π). In particular, the displacement for V makes sense, since at its initial and terminal pieces the argument is constant. Compare V to the curve V' that in the unfolding is a straight line segment from $(0, -1/2)$ to $(2n + 1, 2n + 1/2)$. In the square, V' goes from the lower side to the right one, to the upper one, to the left one, to the lower one, etc., and it reflects from each side at its midpoint. Moreover, the distances of the endpoints of V from the corresponding endpoints of V' are $1/2$. Therefore, when we deform linearly V to V' , we do not cross any point of \mathbb{Z}^2 . This means that the difference of the displacements along those trajectories in the square is less than 2 (this difference may

occur because they end at different points). The displacement along V' is $n + 1/2$, so the displacement along V is between $n - 3/2$ and $n + 5/2$.

The length of V is between the length of V' minus 1 and the length of V' , that is, between $(2n + 1)\sqrt{2} - 1$ and $(2n + 1)\sqrt{2}$. Therefore, as n goes to infinity, the rotation number of V goes to $n/(2n\sqrt{2}) = \sqrt{2}/4$.

If we fix $\varepsilon > 0$ then there is n such that the rotation number of V is larger than $\sqrt{2}/4 - \varepsilon/4$. Then we can choose $\delta > 0$ such that if the obstacle is contained in the disk centered at the origin and diameter less than δ then $(2n + 1, 2n)$ is a vertex of the graph G and any trajectory piece T_n that in the unfolding is a straight line segment from a point of O_0 to a point of $O_{(2n+1, 2n)}$ has rotation number differing from the rotation number of V by less than $\varepsilon/4$ (when we deform linearly V to get this trajectory piece, we do not cross any point of \mathbb{Z}^2). Hence, this rotation number is larger than $\sqrt{2}/4 - \varepsilon/2$.

Now we construct a periodic orbit of admissible type with the rotation number differing from the rotation number of T_n by less than $\varepsilon/2$. By Lemma 2.10, there is a loop A_n in G , passing through $v_n = (2n + 1, 2n)$ and at most 2 other vertices, both from U . As n goes to infinity, then clearly the ratios of displacements and of lengths of T_n and A_n go to 1. Therefore the ratio of their rotation numbers also goes to 1, and if n is large enough, the difference between them will be smaller than $\varepsilon/2$. This gives us δ such that if the obstacle is contained in the disk centered at the origin and diameter less than δ then the set AR contains the number $v > \sqrt{2}/4 - \varepsilon$.

Since AR is symmetric with respect to 0, it contains also the number $-v$, and since it is connected, it contains the interval $[-\sqrt{2}/4 + \varepsilon, \sqrt{2}/4 - \varepsilon]$. \square

Theorem 4.15 also has its counterpart for the billiard in the square.

Theorem 5.8. *For a billiard in a square with a small obstacle, the admissible rotation set is contained in the open interval $(-\sqrt{2}/4, \sqrt{2}/4)$. In particular, $AR \neq R$.*

Since the proof of this theorem utilizes a construction introduced in the proof of Theorem 6.8, we postpone it until the end of Sect. 6.

6. Results on the Full Rotation Set

In this section we will prove several results on the full rotation set R in both cases, not only about the admissible rotation set. Some of the proofs apply to a much more general situation than billiards, and then we will work under fairly general assumptions.

Let X be a compact metric space and let Φ be a *continuous semiflow* on X . That is, $\Phi : [0, \infty) \times X \rightarrow X$ is a continuous map such that $\Phi(0, x) = x$ and $\Phi(s + t, x) = \Phi(t, \Phi(s, x))$ for every $x \in X, s, t \in [0, \infty)$. We will often write $\Phi^t(x)$ instead of $\Phi(t, x)$. Let ξ be an *time-Lipschitz continuous observable cocycle* for (X, Φ) with values in \mathbb{R}^m , that is, a continuous function $\xi : [0, \infty) \times X \rightarrow \mathbb{R}^m$ such that $\xi(s + t, x) = \xi(s, \Phi^t(x)) + \xi(t, x)$ and $\|\xi(t, x)\| \leq Lt$ for some constant L independent of t and x .

The *rotation set* R of (X, Φ, ξ) is the set of all limits

$$\lim_{n \rightarrow \infty} \frac{\xi(t_n, x_n)}{t_n}, \text{ where } \lim_{n \rightarrow \infty} t_n = \infty.$$

By the definition, R is closed, and is contained in the closed ball in \mathbb{R}^m , centered at the origin, of radius L . In particular, R is compact.

Theorem 6.1. *The rotation set R of a continuous semiflow Φ on a connected space X with a time-Lipschitz continuous observable cocycle ξ is connected.*

Proof. Set

$$\psi(t, x) = \frac{\xi(t, x)}{t}$$

for $t > 0$ and $x \in X$. Then the function ψ is continuous on the space $(0, \infty) \times X$. For $n \geq 1$, set $K_n = \psi([n, \infty) \times X)$. With this notation, we have

$$R = \bigcap_{n=1}^{\infty} \overline{K_n}.$$

The set $[n, \infty) \times X$ is connected, so K_n is connected, so $\overline{K_n}$ is connected. Moreover, K_n is contained in the closed ball in \mathbb{R}^m , centered at the origin, of radius L . Therefore $\overline{K_n}$ is compact. Thus, $(\overline{K_n})_{n=1}^{\infty}$ is a descending sequence of compact connected sets, and so its intersection R is also connected. \square

In the case of a billiard on a torus that we are considering, the phase space X is the product of the torus minus the interior of the obstacles with the unit sphere in \mathbb{R}^m (velocities). At the boundaries of the obstacles we glue together the pre-collision and post-collision velocity vectors. This space is compact, connected, and our semiflow (even a flow, since we can move backwards in time, too) is continuous. The observable cocycle is the displacement function. Clearly, it is time-Lipschitz with the constant $L = 1$ and continuous. Thus, Theorem 6.1 applies, and the rotation set R is connected.

Similar situation occurs in the square. Here there is one more complication, due to the fact that there are trajectories passing through vertices. The gluing rule at a vertex q of the square is that the phase points (q, v) and $(q, -v)$ must be identified for all relevant velocities v . Then the flow is also continuous in this case, so Theorem 6.1 also applies. It means that the rotation set R is a closed interval, symmetric with respect to 0.

When we work with invariant measures, we have to use a slightly different formalism. Namely, the observable cocycle ξ has to be the integral of the *observable function* ζ along an orbit piece. That is, $\zeta : X \rightarrow \mathbb{R}^m$ is a bounded Borel function, integrable along the orbits, and

$$\xi(t, x) = \int_0^t \zeta(\Phi^s(x)) ds.$$

Assume that Φ is a continuous flow. Then, if μ is a probability measure, invariant and ergodic with respect to Φ , then by the Ergodic Theorem, for μ -almost every point $x \in X$ the *rotation vector*

$$\lim_{t \rightarrow \infty} \frac{\xi(t, x)}{t}$$

of x exists and is equal to $\int_X \zeta(x) d\mu(x)$.

Problems may arise if we want to use weak- $*$ convergence of measures. If ζ is continuous and μ_n weak- $*$ converge to μ then the integrals of ζ with respect to μ_n converge to the integral of ζ with respect to μ . However, for a general ζ this is not true. Note that in the cases of billiards that we are considering, ζ is the velocity vector. It has a discontinuity at every point where a reflection occurs (formally speaking, it is even not

well defined at those points; for definiteness we may define it there in any way so that it remains bounded and Borel). However, it is well known that the convergence of integrals still holds if the set of discontinuity points of ζ has μ -measure zero (as a random reference, we can give [3], Theorem 7.7.10, p. 234).

Let us call an observable *almost continuous* if the set of its discontinuity points has measure zero for every Φ -invariant probability measure. By what we said above, the following lemma holds.

Lemma 6.2. *If probability measures μ_n weak- $*$ converge to a Φ -invariant probability measure μ and ζ is almost continuous then*

$$\lim_{n \rightarrow \infty} \int_X \zeta(x) d\mu_n(x) = \int_X \zeta(x) d\mu(x).$$

We have to show that this lemma is relevant for billiards.

Lemma 6.3. *Let (Φ, X) be a billiard flow in the phase space. Then the velocity observable function ζ is almost continuous,*

Proof. The only points of discontinuity of ζ are on the boundary of the region Ω in which we consider the billiard. Take a small piece Y of this set. Then for a small $t \geq 0$ the sets $\Phi(t, Y)$ will be pairwise disjoint (if for $y_1 \in \Phi(t_1, Y)$ and $y_2 \in \Phi(t_2, Y)$ with $t_1 \neq t_2$ the velocity is the same, the points of Ω are different). However, by the invariance of μ , their measures are the same. Since the parameter t varies in an uncountable set, those measures must be 0. \square

The following theorem is an analogue of Theorem 2.4 from [7]. Its proof is basically the same as there (except that here we deal with a flow and a discontinuous observable), so we will omit some details. A point of a convex set A is an *extreme* point of A if it is not an interior point of any straight line segment contained in A .

Theorem 6.4. *Let (Φ, X) be a continuous flow and let $\zeta: X \rightarrow \mathbb{R}^m$ be an almost continuous observable function. Let R be the rotation set of (Φ, X, ζ) . Then for any extreme vector \mathbf{u} of the convex hull of R there is a Φ -invariant ergodic probability measure μ such that $\int_X \zeta(x) d\mu(x) = \mathbf{u}$.*

Proof. There is a sequence of trajectory pieces such that the average displacements on those pieces converge to \mathbf{u} . We can find a subsequence of this sequence such that the measures equidistributed on those pieces weakly- $*$ converge to some probability measure ν . This measure is automatically invariant. Therefore, by Lemma 6.2, we can pass to the limit with the integrals of ζ , and we get $\int_X \zeta(x) d\nu(x) = \mathbf{u}$. We decompose ν into ergodic components, and since \mathbf{u} is an extreme point of the convex hull of R , for almost all ergodic components μ of ν we have $\int_X \zeta(x) d\mu(x) = \mathbf{u}$. \square

By Lemma 6.3, we can apply the above theorem to our billiards. In particular, by the Ergodic Theorem, for any extreme vector \mathbf{u} of the convex hull of R there is a point with the rotation vector \mathbf{u} .

Now we will look closer at billiards on the torus \mathbb{T}^m with one obstacle (not necessarily small). We know that its rotation set R is contained in the unit ball. It turns out that although (by Corollary 4.14) R can fill up almost the whole unit ball, still it cannot reach the unit sphere S^{m-1} on a big set. Let us start with the following theorem.

Theorem 6.5. *For a billiard on the torus \mathbb{T}^m with one obstacle, if \mathbf{u} is a rotation vector of norm 1, then there is a full trajectory in \mathbb{R}^m which is a straight line of direction \mathbf{u} .*

Proof. Clearly, such \mathbf{u} is an extreme point of the convex hull of R . By Theorem 6.4, there is an ergodic measure μ such that the integral of the velocity with respect to μ is \mathbf{u} . Thus, the support of μ is contained in the set of points of the phase space for which the vector component is \mathbf{u} . Take t which is smaller than the distance between any two obstacles in the lifting. Then μ -almost all full trajectories of the billiard have direction \mathbf{u} at all times kt , for any integer k . Such a trajectory has to be a straight line with direction \mathbf{u} . \square

The following lemma has been proved in [10] as Lemma A.2.2.

Lemma 6.6. *For every dimension $m > 1$ and every number $r > 0$ there are finitely many nonzero vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in \mathbb{Z}^m$ such that whenever a straight line L in \mathbb{R}^m is at least at the distance r from \mathbb{Z}^m , then L is orthogonal to at least one of the vectors \mathbf{x}_i . In other words, L is parallel to the orthocomplement (lattice) hyperplane $H_i = (\mathbf{x}_i)^\perp$.*

As an immediate consequence of Theorem 6.5 and Lemma 6.6, we get the following result.

Theorem 6.7. *For a billiard on the torus \mathbb{T}^m with one obstacle, the intersection $R \cap S^{m-1}$ is contained in the union of finitely many great hyperspheres of S^{m-1} . The hyperplanes defining these great hyperspheres can be taken as in Lemma 6.6.*

For the billiard in the square with one obstacle, we can determine the full rotation set much better than for the torus case. In the theorem below we do not need to assume that the obstacle is small or even convex. However, we assume that it is contained in the interior of the square and that its boundary is smooth.

Theorem 6.8. *For a billiard with one obstacle in a square, the rotation set is contained in the interval $[-\sqrt{2}/4, \sqrt{2}/4]$.*

Proof. Let Y be the square minus the obstacle. In order to measure the displacement along a trajectory piece T , we have to trace how its lifting to the universal covering space of Y behaves. Since Y is homeomorphic to an annulus, this universal covering has a natural structure of a strip in the plane. Without any loss of generality, we may assume that the displacement along T is positive.

We divide Y into 4 regions, as in Fig. 6.1. The line dividing regions 1 and 2 is a segment of the lowest horizontal straight line such that the whole obstacle is below it; this segment has only its left endpoint belonging to the obstacle. The other three dividing lines are chosen in the same way after turning the whole picture by 90, 180 and 270 degrees. Note that here we are not interested very much to which region the points of the division lines belong, so it is a partition modulo its boundaries.

In the universal covering of Y , our four regions become infinitely many ones, and they are ordered as

$$\dots, 1_{-1}, 2_{-1}, 3_{-1}, 4_{-1}, 1_0, 2_0, 3_0, 4_0, 1_1, 2_1, 3_1, 4_1, \dots \tag{6.1}$$

Here the main number shows to which region in Y our region in the lifting projects, and the subscript indicates the branch of the argument (as in the definition of the displacement) that we are using. Thus, if in the lifting the trajectory goes from, say, 20




Fig. 6.1. Four regions

to 1_{23} , then the displacement is 22 , up to an additive constant. This constant does not depend on the trajectory piece we consider, so it disappears when we pass to the limit to determine the rotation set.

Since we assumed that the displacement along T is positive, then in general, the trajectory moves in the order as in (6.1), although of course it can go back and forth. Look at some region, say, 1_n , which is (in the universal covering) between the region where T begins and the region where T ends. After T leaves 1_n , for good, it can bounce between the left and right sides several times. As this happens, the y -coordinate on the trajectory must grow, so at some point the trajectory will hit the upper side of the square for the first time after it leaves 1_n for good (unless it ends before it does this). We denote the time of this collision by $t(1_n)$. Then we use analogous notation for the time of the first hit of the left side after leaving 2_n for good, etc.

After the trajectory hits the top side at $t(1_n)$, it moves to the left or right (we mean the horizontal component of the velocity). It cannot move vertically, because then it would return to the region 1_n . If it is moving to the left, it is still in the region 2_n , or it just left it, but did not hit the left side of the square yet. Therefore $t(1_n) < t(2_n)$. If it is moving to the right, it is in, or it will return to, the region 2_n . Therefore also in this case $t(1_n) < t(2_n)$.

In such a way we get an increasing sequence of times when the trajectory T hits the consecutive sides of the square (in the lifting). By joining those consecutive reflection points by segments, we get a piecewise linear curve γ , which is not longer than T , but the displacement along γ differs from the displacement along T at most by a constant independent of the choice of T . This curve γ goes from the right side of the square to the upper one, to the left one, to the lower one, to the right one, etc. This is exactly the same behavior that is displayed by the trajectory Γ in the square without an obstacle, that starts at the midpoint of the right side and goes in the direction of $(-1, 1)$. Therefore γ and Γ pass through the same squares in the unfolding. We terminate Γ in that square in the unfolding in which γ ends. Since in the unfolding Γ is a segment of a straight line, it is shorter than γ , again up to a constant independent of γ , and those two curves have the same displacement (as always, up to a constant). This shows that the rotation number of the trajectory piece T is not larger than for the curve Γ , plus a constant that goes to 0 as the length of T goes to infinity. Since the rotation number of Γ (let us think now about the infinite trajectory) is $\sqrt{2}/4$, the limit in the definition of the rotation set cannot exceed this number. \square

The two trajectories in the square without an obstacle, described in the last paragraph of the above proof, are also trajectories of any square billiard with one small obstacle.

Therefore in this case the rotation set R contains the interval $[-\sqrt{2}/4, \sqrt{2}/4]$. In such a way we get the following result.

Theorem 6.9. *For a billiard with one small obstacle in a square, the rotation set is equal to the interval $[-\sqrt{2}/4, \sqrt{2}/4]$.*

Now we present the proof of Theorem 5.8 that has been postponed until now.

Proof of Theorem 5.8. Let us use the construction from the proof of Theorem 6.8 and look at a long trajectory piece T of admissible type. We get an increasing sequence of times when the trajectory T hits the consecutive sides of the square (in the lifting). Construct a partial unfolding of T , passing to a neighboring square only at those times. In such a way we get a piecewise linear curve T' which sometimes reflects from the sides of the unfolded square, and sometimes goes through them. Its length is the same as the length of T . Moreover, the curve γ , constructed in the proof of Theorem 6.8 starts and terminates in the same squares as T' and as we know from that proof, the displacements along γ and T differ at most by a constant independent of T . It is also clear that the same holds if we replace γ by the segment Γ (also from the proof of Theorem 6.8). Thus, up to a constant that goes to 0 as the length of T goes to infinity, the rotation number of T is $\sqrt{2}/4$ multiplied by the length of Γ and divided by the length of T' .

By the same reasons as in the proof of Theorem 4.15, there exist positive constants $c_1 < c_2$ and $\alpha > 0$, such that for every trajectory piece of admissible type the distance between two consecutive reflections from obstacles is contained in $[c_1, c_2]$, and the direction of a trajectory piece of admissible type changes by at least α at each reflection. However, we have to take into account that the direction of T' can change also at the reflections from the boundaries of the squares, and then we do not know how the angle changes. Thus, either immediately before or immediately after each reflection from an obstacle there must be a piece of T' where the direction differs from the direction of Γ by at least $\alpha/2$. Such a piece has length larger than or equal to the distance from the obstacle to the boundary of the square, which is at least $c'_1 = 1/2 - \sqrt{2}/4$. Those pieces are alternating with the pieces of T' of length at most c_2 each, where we do not know what happens with the direction. This means that as we follow T' (except the initial piece of the length bounded by c_2), we move in the direction that differs from the direction of Γ by at least $\alpha/2$ for at least $c'_1/(c'_1 + c_2)$ of time. Therefore, in the limit as the length of T goes to infinity, the length of Γ divided by the length of T' is not larger than

$$b = \frac{c_2}{c'_1 + c_2} + \frac{c'_1}{c'_1 + c_2} \cos \frac{\alpha}{2}.$$

This proves that $AR \subset [-b\sqrt{2}/4, b\sqrt{2}/4]$. Since $b < 1$, this completes the proof. \square

References

1. Alesdà, L., Llibre, J., Misiurewicz, M.: Combinatorial dynamics and entropy in dimension one. Advanced Series in Nonlinear Dynamics, Vol. 5, Second Edition, River Edge, NJ: World Scientific, 2000
2. Bárány, I., Simányi, N.: A note on the size of the largest ball inside a convex polytope. Period. Math. Hungar. 51, 15–18 (2005)
3. Bauer, H.: Probability Theory and Elements of Measure Theory. New York: Holt, Reinhart and Winston, 1972

4. Chernov, N.: Construction of transverse fiberings in multidimensional semidispersed billiards. *Funct. Anal. Appl.* **16**, 270–280 (1982)
5. Dettmann, C. P.: The Lorentz Gas: A Paradigm for Nonequilibrium Stationary States. In *Hard Ball Systems and the Lorentz Gas*, (ed. D. Szász), *Encyclopaedia Math. Sci.* **101**, Berlin-Heidelberg-New York: Springer Verlag, 2000, pp. 315–365
6. Katok, A., Hasselblatt, B.: *Introduction to the Modern Theory of Dynamical Systems*. Cambridge: Cambridge University Press, 1995
7. Misiurewicz, M., Ziemian, K.: Rotation sets for maps of tori. *J. London Math. Soc. (2)* **40**, 490–506 (1989)
8. Misiurewicz, M., Ziemian, K.: Rotation sets and ergodic measures for torus homeomorphisms. *Fund. Math.* **137**, 45–52 (1991)
9. Stoyanov, L.: An estimate from above of the number of periodic orbits for semi-dispersed billiards. *Commun. Math. Phys.* **124**, 217–227 (1989)
10. Szász, D.: The K-property of ‘Orthogonal’ Cylindric Billiards. *Commun. Math. Phys.* **160**, 581–597 (1994)
11. Ziemian, K.: Rotation sets for subshifts of finite type. *Fund. Math.* **146**, 189–201 (1995)

Communicated by G. Gallavotti