

ON DYNAMICS OF VERTICES OF LOCALLY CONNECTED POLYNOMIAL JULIA SETS

A. BLOKH AND G. LEVIN

December 8, 2000; revised version August 7, 2001; new version January 4, 2002

ABSTRACT. Let P be a polynomial whose Julia set J is locally connected. Then a non-preperiodic non-precritical vertex of J must have the limit set which coincides with the limit set of an appropriately chosen recurrent critical point of P . In particular, if all critical points of P are non-recurrent then all vertices of J are preperiodic or precritical.

1. INTRODUCTION

In this paper P denotes a complex polynomial with the Julia set J . Properties of $P|_J$ and topological properties of J have been studied in holomorphic dynamics for quite some time (see e.g. [BH], [DH], [Do], [F], [H], [J], [McM], [Mi], [Sul], [Th]). From now on we assume in this paper that J is locally connected and study the orbits of the vertices of J . As a corollary we prove that if all critical points of P are non-recurrent then all vertices of J are preperiodic or precritical.

It is known that if J is locally connected then it is connected. By a Caratheodory theorem [CL], any point $x \in J$ is *accessible* from the basin of attraction of infinity A_∞ by one or several paths tending to x which can be chosen to be so-called *external rays* [DH]. Let $N(x)$ be the number of such rays, and call points x with $N(x) \geq 3$ *vertices* of J (one can also define $N(x)$ as the number of components of the set $J \setminus \{x\}$). In the language of continuum theory, vertices are *cut points* cutting the Julia set in at least three components. The following problem is fundamental for understanding the topology of Julia sets; it was posed in the framework of polynomial laminations and solved for quadratic laminations in [Th].

Problem 1. *Is it true that all vertices of the locally connected Julia set are either pre-critical or preperiodic?*

If we think of J as a “graph” with, perhaps, infinitely many vertices then it is natural to expect that in some cases P acts on J similarly to the way it would act on finite “graphs”. Now, if a finite “graph” G is invariant for P then all vertices of G are precritical

2000 *Mathematics Subject Classification* Primary 37F10; Secondary 37E25.

Key words and phrases. Julia set, vertices, laminations, recurrent critical points.

The first author was partially supported by NSF grant DMS 9970363.

or preperiodic (a point is *preperiodic* - resp. *precritical* - if it is mapped onto a periodic - resp. critical - point by $f^k, k \geq 0$). This makes the question above rather natural.

Related questions in the degree higher than 2 were considered by Kiwi [Ki] who proved that for a polynomial of degree d the number $N(x)$ of rays landing at a non-preperiodic non-precritical point $x \in J$ is at most d (this is another proof of the result of [Th] in the quadratic case). In [BL1, BL2] we obtained further results one of which is described in Theorem 2; it is easy to see, that it implies the results mentioned above.

By the *grand orbit* (of x) we mean the union of all preimages of all iterates of x . Note that the number $N(x')$ is the same for all points x' of the grand orbit of x if this orbit contains no critical points. By a *Siegel disc* we understand a Fatou component whose boundary is a simple closed curve such that an appropriate power of the map P on its boundary is an irrational rotation.

Theorem 2 [BL1, BL2]. $\sum_{x \in \Gamma} (N(x) - 2) \leq k_{\sim} - k_S - 1 \leq d - 2$, where Γ is a non-empty collection of non-preperiodic non-precritical points $x \in J$ from distinct grand orbits with $N(x) \geq 3$, k_{\sim} is the number of distinct grand orbits of non-preperiodic critical points $c \in J(P)$ with $N(P(c)) = 1$ (i.e., $P(c)$ is the landing point of exactly one external ray) and k_S is the number of periodic orbits of the Siegel discs.

In fact, a complete analog of Theorem 2 is proven in [BL1, BL2] for invariant *laminations* and their factors which implies (as a particular case) Theorem 2 for polynomials with locally connected Julia sets.

In a way, Theorem 2 already shows to what extent the dynamics of a polynomial on its locally connected Julia set is similar to the dynamics of a polynomial on a finite graph: even if non-preperiodic non-precritical vertices exist, they form at most $d - 2$ grand orbits. Observe, that although in general Theorem 2 does not solve Problem 1 and can only be considered as a step on the way to its solution, in some particular cases it implies that there are no non-preperiodic non-precritical vertices of a locally connected Julia set. In particular, Theorem 2 implies the following corollary.

Corollary 3 (cf. [Do], [Po]). *Let P be a polynomial whose critical points are attracted by a periodic orbit or preperiodic. Then all vertices of J are preperiodic or precritical.*

In the present paper we suggest another class of polynomials such that all non-precritical vertices of their Julia sets are preperiodic. A polynomial P is called *semi-hyperbolic* [CJY] if no critical point of P in J is recurrent and there are no parabolic cycles. As follows from [CJY], the Julia set of such polynomial is locally connected if it is connected, and hence every $x \in J$ is the landing point of a finitely many external rays ([Ki], [BL1], [BL2]). Assume in addition that P has only repelling periodic points (and so its Julia set is a *dendrite*, i.e. locally connected tree-like continuum). Then using Yoccoz puzzle structure for P we prove that all vertices of J are preperiodic or precritical (we briefly outline the proof in Section 2). In fact, the same result holds for a wider class \mathcal{A} of all polynomials with no recurrent critical points and with locally connected Julia set. Thus, compared to the above case in \mathcal{A} we allow attracting or parabolic cycles.

If J is a locally connected Julia set then $x \in J$ is a *local vertex* of J if there is a triod $Y \subset J$ whose branch point is x . If J is a dendrite, this is equivalent to the usual definition, yet if there are bounded Fatou domains, the set of local vertices may be wider.

Main Theorem. *If $P \in \mathcal{A}$ then every local vertex of J is either preperiodic or precritical.*

A complication for $P \in \mathcal{A}$ is that J no longer is a dendrite; we deal with it by developing topological dynamics of *piecewise local homeomorphisms* of *unshielded* plane continua which are *backward stable* (see definitions in Section 4), more precisely by describing the limit sets of local vertices of such maps. This description, applied to laminations and polynomials, yields Theorem 13 below which in turns implies the Main Theorem.

Theorem 13. *Suppose that one of the assumptions below holds:*

- (1) *\sim is an invariant lamination and $f : J \rightarrow J$ is the corresponding factor map;*
- (2) *f is a polynomial, J is its Julia set, and J is locally connected.*

Then if x is a non-preperiodic non-precritical local vertex of J then $\omega(x)$ coincides with $\omega(c(x))$ for a recurrent critical point $c(x)$ depending on x .

The paper is organized as follows. In Section 2 we outline the proof of Main Theorem for semi-hyperbolic polynomials. In Section 3 we introduce the notion of *unshielded* continua and study their topology (connected Julia sets are unshielded continua). In Section 4 we study backward stable continuous maps of unshielded continua which are local homeomorphisms outside a finite set and prove Theorem 13.

We want to thank L. Oversteegen for useful discussions of the proofs of Lemmas 5 and 6 and to M. Misiurewicz for valuable discussions of the proof of Theorem 11. Also, we want to thank the referee for finding a gap in the original proof of the Main Theorem and one of the lemmas and for making valuable remarks which helped us improve the style of this paper.

2. SEMI-HYPERBOLIC POLYNOMIALS WITHOUT ATTRACTING CYCLES

In this section we consider a special case of semi-hyperbolic polynomials with all periodic points repelling, and outline the idea of proving Main Theorem in this case.

Let P be a polynomial of degree $d > 1$ whose Julia set J is connected and periodic points repelling. Assume that P is a *semi-hyperbolic* polynomial [CJY] (i.e. every critical point in J is not recurrent). By [CJY], the Julia set of such polynomial is locally connected, and hence, every $x \in J$ is a landing point of a finitely many external rays.

To demonstrate the idea of the proof in this case, consider Yoccoz puzzle structure for P . Note that in our case P is not renormalizable, and the puzzle structure is constructed in a standard way (see e.g. [H], [BL1]). Recall that for every $n = 0, 1, \dots$ there are finitely many pairwise disjoint open puzzle pieces $Y_n^1, \dots, Y_n^{k_n}$ of the level n . For any $Y = Y_n^i$, the intersection of the boundary ∂Y of Y with the Julia set has finitely many points (preperiodic by P). The union of the closures of $Y_n^i, 1 \leq i \leq k_n$ contains the Julia set J , and P maps any puzzle piece of the level $n > 0$ onto a puzzle piece of the level $n - 1$.

Given a point $z \in J$, let $Y_n(z)$ be a nested sequence of open puzzle pieces, such that $z \in \overline{Y_n(z)}$ for any n . In [L], [BL1] we show that the Julia set J is locally connected if and only if the intersection of all $Y_n(z)$ is the single point z , for each $z \in J$.

Given a puzzle piece $Y = Y_n(z)$, we define a finite tree $T_n(z)$ as follows. As we know, the intersection of the boundary ∂Y of Y with the Julia set consists of a finitely many points x_1, \dots, x_p . Given any two of them x_i, x_j , let $[x_i, x_j]$ be the unique arc in J with

the ends x_i, x_j (such an arc exists because J is locally connected). Define now $T_n(z)$ as the union of all possible arcs $[x_i, x_j]$. It is easy to see that $T_n(z)$ is a finite tree.

Then we use the fact that every critical point is non-recurrent and rely upon a result of [BM] to find a number m and a sequence n_k such that the map P^{n_k} maps homeomorphically the piece $Y_{m+n_k}(z)$ onto a piece of a fixed level m . Since there are finitely many pieces on each level, one can assume that $P^{n_k}(Y_{m+n_k}(z))$ is a fixed piece Y_m .

Let us now assume that z is neither preperiodic nor precritical, and prove that then z is the landing point of at most 2 external rays. Assume (by contradiction) that there are at least 3 external rays landing at z . Then, for every n large enough, z is a vertex of every tree $T_n(z)$. From the above, P^{n_k} maps $T_{n_k+m}(z)$ homeomorphically onto some tree t_k , and $z_k = P^{n_k}(z)$ is a vertex of t_k . By the construction, t_k is a subset of J , and each end point of t_k is a point of the set $\partial Y_m \cap J$. Since the latter set is finite, and Y_m is a fixed piece, there are finitely many trees t_k , each of which is finite. Therefore, the points z_k belong to a finite set of the vertices of the trees t_k . Thus $z_k = z_i$ for different k, i , which means that the point z is preperiodic.

3. TOPOLOGICAL PROPERTIES OF LOCALLY CONNECTED UNSHIELDED CONTINUA

A plane continuum K is said to be *unshielded* [BO] if it coincides with the boundary of a domain U which is a connected component of the complement of K . Connected Julia sets are unshielded since they coincide with the boundary of A_∞ . In this section we study topological properties of locally connected unshielded continua.

It immediately follows from the definition that an unshielded continuum K cannot contain a continuum of the shape of the letter θ . Therefore, the following so-called θ -curve Theorem applies to unshielded continua.

Theorem 4 (θ -curve Theorem) [Kur]. *A locally connected unshielded continuum is hereditarily locally connected: all its subcontinua are locally connected. Also, any connected subset of a locally connected unshielded continuum is arcwise connected.*

We need some standard definitions. An *arc* is a homeomorphic image of the interval $[0, 1]$. If the endpoints of an arc are a, b and if it causes no confusion we use the notation $[a, b]$.

The following lemma is proven in [BO] and is given here for the sake of completeness.

Lemma 5 [BO]. *Let T be a locally connected unshielded continuum. For any sequence of continua $M_n \subset J$ such that diameters of sets M_n do not converge to 0 there exists a subsequence K'_n such that all sets K'_n contain some arc L' .*

Proof. We can always find a subsequence of continua M_n converging in the Hausdorff sense to a non-degenerate continuum. By Theorem 4 all M_n are locally connected and therefore arcwise connected. Choose subarcs K_n of M_n which contain points $u_n, v_n \in K_n$ positively distant from each other. We can assume that there are points $x \neq y$ such that $u_n \rightarrow x, v_n \rightarrow y$. Assuming that u_n and v_n are very close to x, y respectively, and using the fact that T is locally connected, we can find small disjoint arcs $L'_n = [x, u_n] \subset T, R'_n = [y, v_n] \subset T$ so that $\lim L'_n = \{x\}$ and $\lim R'_n = \{y\}$

Since K_n is arcwise connected by Theorem 4, we can always find an arc Q_n connecting u_n and v_n inside K_n . Then we can always choose points $x_n \in Q_n \cap L'_n$ and $y_n \in R'_n \cap Q_n$

so that the subarc $[x_n, y_n]$ of Q_n connecting x_n and y_n inside Q_n is disjoint from $L'_n \cup R'_n$ outside its endpoints x_n, y_n . Let $L_n = [x, x_n] \subset L'_n$ ($R_n = [y, y_n] \subset R'_n$) be the arcs connecting x and x_n (y and y_n) inside L'_n (R'_n). Then $J_n = [x, x_n] \cup [x_n, y_n] \cup [y_n, y]$ is an arc, $\lim L_n = \{x\}$ and $\lim R_n = \{y\}$. We can choose a subsequence now so that $[x_n, y_n]$ converge to the limit continuum K ; observe that since L_n, R_n converge to points, the arcs J_n converge to the same continuum K . Let $X' = K \cup_n J_n$, then $X' \subset T$ is a continuum.

Choose an arc $A' = [x, y] \subset K$ connecting x and y . Then $[x, x_n] \cup [x_n, y_n] \cup [y_n, y] \cup [y, x] = E_n$ is a closed curve for every n . Since L_n and R_n are very small for big n , we may assume that there exists a non-degenerate arc $A \subset A' \setminus \cup_n [L_n \cup R_n]$. If $A \subset J_n$ for all n then $A \subset [x_n, y_n] \subset K_n$ and we are done. Assume that there exists a point $a \in A \setminus J_1$. Then the fact that $E_1 = [x, x_1] \cup [x_1, y_1] \cup [y_1, y] \cup [y, x]$ is a closed curve implies that there exists a bounded region D complementary to E_1 such that a is contained in $\partial D = S$. By Theorem 4 S is locally connected; since $S \subset T$ we see that S is a simple closed curve. Then $A_1 = S \cap A'$ is a non-degenerate arc containing a (both S and A' are contained in an unshielded continuum T , and the intersection of an arc and a simple closed curve in an unshielded continuum is an arc or a point unless it is empty). Let $I_1 = \overline{S \setminus A'}$. Since T contains no θ -curve, any pair of distinct points $p, q \in S$ with $p \in I_1$ and $q \in A_1$ cuts each J_n between x and y . Hence either A_1 or I_1 is contained in infinitely many J_n . Since $L_n \rightarrow \{x\}, R_n \rightarrow \{y\}$ it is easy to find a non-degenerate arc L' which is contained in infinitely many K_n as required. ■

Let us now introduce the following notion: a point x of a hereditarily locally connected continuum K is called a *locally ε -essential vertex* if there exists a triod $Y \subset K$ whose vertex is the point x and whose branches are all of diameters at least ε (in this paper we deal only with locally connected continua and thus give the definition in this particular case). A point which is a locally ε -essential vertex for some $\varepsilon > 0$ is called a *local vertex* of K . Observe that a local vertex does not have to be a vertex (for example, if a continuum T has the shape of a letter σ then the only branching point of T is a local vertex but not a vertex of T). Now, a point x of a hereditarily locally connected continuum X is said to be a *globally ε -essential vertex* if there are at least three components of $X \setminus \{x\}$ of diameter at least ε . Clearly, a globally ε -essential vertex is locally ε -essential, and all globally ε -essential vertices of T taken over all $\varepsilon > 0$ form the set of all vertices of T .

The following topological result is used in the proof of Theorem 10.

Lemma 6. *Let T be a locally connected unshielded continuum. Then for every ε the number of locally ε -essential vertices of T is finite (so in particular, the set of all local vertices is at most countable).*

Proof. Suppose that there are infinitely many locally ε -essential vertices $\{a_i\}_{i=1}^\infty$. For every i let Y_i be a triod containing a_i as its vertex and having branches N_i, L_i and M_i each of which has diameter greater than ε . Assume that $a_i \rightarrow a$. By Lemma 5 we can choose a subsequence and an arc B so that $a \notin B, a_i \notin B$ for any i , and $B \subset N_i$ for any i (indeed, we can always choose a subsequence and B by Lemma 5 so that $a \notin B$ and then refine this subsequence so that $a_n \notin B$ for any n). Then we can refine this subsequence and choose another subsequence and an arc C so that $a \notin C, a_i \notin C$ for any i and $C \subset L_i$ for any i ; clearly then C and B are disjoint. Finally, we can refine the latest subsequence

and find its subsequence and an arc D so that $a \notin D, a_i \notin D$ for any i and $D \subset M_i$ for any i ; clearly then C, B and D are disjoint. Now, Y_1 contains B, C and D and so does Y_2, Y_3, \dots . However, this is clearly impossible for the unshielded continuum T to contain all these triods since in this case it is not unshielded, a contradiction. ■

4. MAIN THEOREM

So far the arguments were purely topological; now it is the time to apply dynamical arguments and prove the Main Theorem. Our plan is to obtain some general results for a certain class of continuous maps of locally connected unshielded continua. Since polynomial maps on their locally connected Julia sets belong to this class of maps we then apply our results and deduce Theorem 10, Theorem 13 and the Main Theorem.

First we need the following definition. A map $f : X \rightarrow X$ of a metric space X into itself is said to be a *local homeomorphism at a point x* if any small neighborhood of x is homeomorphically mapped by f onto a neighborhood of $f(x)$. A map is called a *local homeomorphism on a set $Y \subset X$* if it is a local homeomorphism at every point of Y . We study local homeomorphisms of connected neighborhoods in a locally connected unshielded continuum below.

Lemma 7. *Let $f : T \rightarrow T$ be a continuous map of a locally connected unshielded continuum T . Let $U \subset T$ be an open connected set and V be a component of $f^{-1}(U)$ such that $f|_V$ is a local homeomorphism. Then V maps onto U and, moreover, $f(\partial V) = \partial U$.*

Proof. As a component of an open set in a locally connected space, V is open. Since $f|_V$ is a local homeomorphism, it is an open map, hence $f(V) \subset U$ is open. Let us show that $W = U \setminus f(V)$ is open too. Indeed, if $x \in W$ and yet no neighborhood of x is inside W then x is the limit point of a sequence of points $f(y_n), y_n \in V$. Assume that $y_n \rightarrow y \in \partial V$ (y cannot belong to V because $f(y) \notin f(V)$). Then $V \cup \{y\}$ is connected and maps into U , a contradiction with the choice of V . Hence, W is open. But this together with the connectedness of U implies that W is empty and $f(V) = U$.

Let us show that $f(\partial V) = \partial U$. Indeed, otherwise one of the two cases must take place. First, there may exist a point $x \in \partial V$ which maps inside U . However then $V \cup \{x\}$ is connected and maps into U , a contradiction with the choice of V . Second, if $y \in \partial U$ then because clearly $f(\bar{V}) = \bar{U}$ we see that $y = f(z)$ for some $z \in \bar{V}$, and $z \in V$ is impossible since then $f(z)$ would belong to U . Thus, $z \in \partial V$ as desired. ■

The next lemma deals with local homeomorphisms of sufficiently small connected neighborhoods of local vertices.

Lemma 8. *Let $U \subset T$ be a connected neighborhood of a local vertex x in a locally connected unshielded continuum T and $f : T \rightarrow T$ is a map such that $f|_U$ is a local homeomorphism and U is a component of $f^{-1}(f(U))$. Then the following holds:*

- (1) *for any simple closed curve $S \subset f(U)$ any component R of its preimage inside U is a simple closed curve such that $f(R) = S$;*
- (2) *for any simple closed curve $R \subset U$ the set $f(R)$ is a simple closed curve too;*
- (3) *the images of different components of $U \setminus \{x\}$ under a local homeomorphism $f : U \rightarrow T$ are disjoint.*

Proof. (1) A set R is a locally connected continuum since T is hereditarily locally connected by Theorem 4. Moreover, since f is a local homeomorphism on R then R maps onto S and has no endpoints or vertices, i.e. R is a simple closed curve.

(2) Clearly, $f(R)$ contains a simple closed curve S' , and so there exists a component R' of the preimage of S' having a non-trivial intersection with R . Since T is unshielded, this implies that $R' = R$ and $S' = S$ as desired.

(3) Assume that there are components Y_1, Y_2 of $U \setminus \{x\}$ and points $y_1 \in Y_1, y_2 \in Y_2$ with $f(y_1) = f(y_2) = z$. By Theorem 4 there exist arcs $[x, y_1] \subset Y_1$ and $[x, y_2] \subset Y_2$. Since f is a homeomorphism on a small neighborhood of x we may assume that there are points $z_1 \in (x, y_1)$ and $z_2 \in (x, y_2)$ such that $f(z_1) = f(z_2)$ and images of (x, z_1) and (x, z_2) are disjoint. Then $S = f[x, z_1] \cup f[x, z_2] \subset f(U)$ is a simple closed curve, so by (1) there exists a simple closed curve $R \supset [x, z_1] \cup [x, z_2]$ which contradicts the fact that $(x, z_1]$ and $(x, z_2]$ are contained in distinct components of $U \setminus \{x\}$. ■

If $f : T \rightarrow T$ is a local homeomorphism of a locally connected unshielded continuum T everywhere but at a finite set C_f of *critical* points then we call it a *piecewise local homeomorphism*. Choose a metric $d(\cdot, \cdot)$ on T . For $x \in T$ and $n > 0$ consider the family $\mathcal{B}(x, n)$ of all connected neighborhoods U of $f^n(x)$ which can be pulled back to x as local homeomorphisms. Thus, we include U in $\mathcal{B}(x, n)$ if U is a connected neighborhood of $f^n(x)$ and the component V of $f^{-n}(U)$ which contains x is such that $f^n|_V$ is a local homeomorphism. Then the union of any collection of neighborhoods from $\mathcal{B}(x, n)$ belongs to the same family and there exists the maximal neighborhood $U_n(x) = U \in \mathcal{B}(x, n)$ and the corresponding neighborhood $V_n(x) = V$ of x . Denote the minimal distance $d(f^n(x), \partial U)$ between $f^n(x)$ and ∂U by $r_n(x) = r_n$. Then $r_n > 0$ if all points $x, \dots, f^{n-1}(x)$ are not critical, otherwise define r_n as 0.

By the definition there are points z on the boundary ∂V of V such that f^n is not a local homeomorphism at z and $d(f^n(z), f^n(x)) = r_n$ (otherwise we could extend V and increase r_n). Hence there exists a point $z \in \partial V$, a critical point $c_n(x) = c_n$ and a number $m_n < n$ such that $f^{m_n}(z) = c_n$ and $d(f^n(z), f^n(x)) = r_n$. Let us say that c_n *generates* r_n . Following [BM] (and mimicking the terminology due to Yoccoz) we call points with $r_n \rightarrow 0$ *persistent*, and points with $r_n \not\rightarrow 0$ *reluctant*. For example, preimages of critical points are persistent while repelling periodic non-critical points are reluctant.

Let T be a metric continuum. Say that $f : T \rightarrow T$ is *backward stable* if for any δ there exists ε such that for any continuum K with $\text{diam}(K) \leq \varepsilon$, any $n \geq 0$ and any component M of $f^{-n}(K)$, $\text{diam}(M) \leq \delta$ (such components are called *pull-backs*).

Lemma 9. *Let $f : T \rightarrow T$ be a backward stable piecewise local homeomorphism. Then:*

- (1) *for any continuum K there exists $\varepsilon > 0$ such that $\text{diam}(f^n(K)) \geq \varepsilon$ for all n ;*
- (2) *if the set $\overline{\text{orb}(x)}$ does not contain critical points then x is reluctant.*

Proof. (1) Let $\delta = \text{diam}(K)$. Choose ε so that for any continuum K' with $\text{diam}(K') \leq \varepsilon$, any $n \geq 0$ and any component M of $f^{-n}(K')$, $\text{diam}(M) \leq \delta/2$. Then $\text{diam}(f^n(K)) > \varepsilon$ for any n since otherwise we take the component M of $f^{-n}(f^n(K))$ containing K and get a contradiction because $M \supset K$ and hence with $\text{diam}(M) \geq \delta$.

(2) Let $d(\overline{\text{orb}(x)}, C_f) = \delta$. Choose ε such that if M is a continuum, $\text{diam}(M) < \varepsilon$ then all its pull-backs are of diameter less than δ . For $f^n(x)$ choose its connected neighborhood

V of diameter less than ε so that its boundary is sufficiently far away from $f^n(x)$. Then f^n restricted on the pull-back of V containing x is a local homeomorphism as desired. ■

Fix a locally connected unshielded plane continuum T and consider a class \mathcal{A}_T of continuous *backward stable* maps $f : T \rightarrow T$ of T into itself such that f is a local homeomorphism at all points of T except of a finite set of points called *critical*. Now we prove Theorem 10 describing orbits of reluctant vertices under maps from \mathcal{A}_T .

Theorem 10. *Let $f : T \rightarrow T, f \in \mathcal{A}_T$, be a map of a locally connected unshielded continuum T . If a local vertex of T is reluctant, then it is preperiodic.*

Proof. Suppose that x is a reluctant local vertex. Then there exist $\varepsilon > 0$, a sequence of integers $n_k \rightarrow \infty$ and a sequence of connected neighborhoods U_k of $f^{n_k}(x)$ such that the minimal distance $d(f^{n_k}(x), \partial U_k)$ is greater than ε and components V_k of preimages $f^{-n_k}(U_k)$ containing x are such that $f^{n_k}|_{V_k}$ is a local homeomorphism.

Choose a triod $Y = [y_1, x] \cup [y_2, x] \cup [y_3, x] = Y_1 \cup Y_2 \cup Y_3$ (it is possible since x is a local vertex) and consider a few cases. The simplest is when for infinitely many k the components B_1^k, B_2^k and B_3^k of $V_k \setminus \{x\}$ covering small semi-neighborhoods of x on Y_1, Y_2 and Y_3 , are distinct. Fix one such k and denote the components of $B_i^k \cap Y_i$ by $[z_i, x], 1 \leq i \leq 3$. A point z_i may well coincide with y_i , but if not then z_i must come from the boundary of V_k . By Lemma 8(3), sets $f^{n_k}(z_1, x), f^{n_k}(z_2, x), f^{n_k}(z_3, x)$ are pairwise disjoint. Let us show that they have a bounded away from zero diameters. Indeed, if $z_i = y_i$ this holds by Lemma 9(1) (which implies that the diameters of the images of Y_i are bounded away from zero), and if $z_i \neq y_i$ this follows by Lemma 7 (which yields that $f^{n_k}(z_i) \in \partial U_k$ and $d(f^{n_k}(x), \partial U_k) \geq \varepsilon$). This means that $f^{n_k}(x)$ are locally ε' -essential vertices for some $\varepsilon' > 0$, and so by Lemma 6 the point x is preperiodic.

Now assume that for all k , small semi-neighborhoods of x on Y_1, Y_2 are contained in one component C_k of $V_k \setminus \{x\}$. Since C_k is arcwise connected by Theorem 4, there is an arc $[u_1, u_2] \subset C_k$ with $u_1 \in Y_1 \setminus \{x\}, u_2 \in Y_2 \setminus \{x\}$ and $(u_1, u_2) \cap (Y_1 \cup Y_2) = \emptyset$. Since T is unshielded, no two points on Y_1, Y_2 which are closer to x are connected by a similar arc. Moreover, $S = [u_1, x] \cup [x, u_2] \cup [u_2, u_1]$ is a simple closed curve, and since by the assumptions $x, u_1, u_2 \in V_k$ then (again because T is unshielded) $S \subset V_k$ for any k . By Lemma 9(1), the diameter of $f^{n_k}(S)$ is greater than some ε' for all k , and by Lemma 8(2) the set $f^{n_k}(S)$ is a simple closed curve. Hence we can two arcs I'_k, I''_k in $f^{n_k}(S)$ of diameter $\varepsilon'/3$ which intersect only over their common endpoint $f^{n_k}(x)$.

Now, since T is unshielded, a small semi-neighborhood of x in Y_3 is not contained in C_k , and so there is a point $z_3^k \in Y_3$ such that $[z_3^k, x)$ is contained in a component of $V_k \setminus \{x\}$ distinct from C_k and either $z_3^k \in \partial V_k$ or $z_3^k = y_3$. As before, this implies that the set $f^{n_k}[z_3^k, x)$ has a bounded away from zero diameter and is disjoint from I'_k, I''_k , and again as before this implies that x is preperiodic. ■

Let us prove Theorem 11 which deals with the behavior of persistent vertices of a locally connected unshielded continuum T under a map $f \in \mathcal{A}_T$. Observe that the ideas we use in the proof are quite similar to the ones used in [BM] where we deal with more general situation under weaker than backward stability assumptions about the map.

Theorem 11 (cf. [BM]). *Let $f : T \rightarrow T, f \in \mathcal{A}_T$ be a map of a locally connected unshielded continuum T . If a point x of T is persistent then either it is precritical or its*

limit set coincides with the limit set of a recurrent critical point $c(x)$ depending on x .

Proof. Let x be a non-precritical persistent point. For every n find a connected neighborhood V of x , a connected neighborhood U of $f^n(x)$, a critical point c_n and a number m_n such that $f^n(V) = U$, $f^n|_V$ is a local homeomorphism, $c_n \in \partial f^{m_n}(V)$, $f^{n-m_n}(c_n) \in \partial U$ and $d(f^{n-m_n}(c_n), f^n(x)) = r_n$. Let us show that then the maximum of distances $d(c_n, f^{m_n}(x)), d(f(c_n), f^{m_n+1}(x)), \dots, d(f^{n-m_n}(c_n), f^n(x))$ converges to 0 as $n \rightarrow \infty$.

Indeed, given $\varepsilon' > 0$ choose ε so that for any continuum $K \subset T$ with $\text{diam}(K) \leq \varepsilon$, the diameter of any component of $f^{-l}(K)$ is less than ε' , for all $l \geq 0$. Now, cover T with finitely many open connected neighborhoods of diameter less than ε (it is possible since T is locally connected). Let δ_ε be the Lebesgue number for this cover. This means that if $r_n < \delta_\varepsilon$ then for the point $f^n(x) \in T$ there exists a connected neighborhood from our cover U of diameter less than ε such that the ball of radius r_n centered at $f^n(x)$ is contained in U . In particular, $f^n(x)$ and $f^{n-m_n}(c_n)$ belong to the same connected neighborhood U of diameter less than ε . By the choice of ε this in turn implies that sets $f^{m_n}(\bar{V}), f^{m_n-1}(\bar{V}), \dots, f^n(\bar{V}) = \bar{U}$ have diameters at most ε' . Since $c_n, f^{m_n}(x) \in f^{m_n}(\bar{V}); f(c_n), f^{m_n+1}(x) \in f^{m_n+1}(V); \dots; f^{n-m_n}(c_n), f^n(x) \in U$ we can easily see that the maximum of distances $d(c_n, f^{m_n}(x)), d(f(c_n), f^{m_n+1}(x)), \dots, d(f^{n-m_n}(c_n), f^n(x))$ is less than ε' which proves that in fact this maximum converges to 0 as $n \rightarrow \infty$.

In short, this means the following. Let C be a finite collection of critical points of f . For any n we choose a number m_n and a point $c_n \in C$ such that the orbit segment $c_n, f(c_n), \dots, f^{n-m_n}(c_n)$ approximates the orbit segment $f^{m_n}(x), \dots, f^n(x)$ with the error of approximation s_n (meaning that the distance $d(f^i(c_n), f^{m_n+i}(x)), 0 \leq i \leq n - m_n$ is at most s_n) and $s_n \rightarrow 0$ as $n \rightarrow \infty$.

Let us show that this implies that there exists a recurrent point $t \in C$ such that $\omega(t) = \omega(x)$. First of all, let Q be the set of those points $q \in C$ which appear as points c_n infinitely many times. Then the fact that $s_n \rightarrow 0$ implies that $Q \subset \omega(x)$. Indeed, if $q \in Q$ then $d(f^{m_{n_k}}(x), q) \leq s_{n_k} \rightarrow 0$ along a sequence $n_k \rightarrow \infty$. Moreover, since x by the assumption is never mapped into a critical point then $m_{n_k} \rightarrow \infty$ as well. Hence $q \in \omega(x)$. Therefore $\omega(Q) = \cup_{q \in Q} \omega(q) \subset \omega(x)$.

On the other hand, $d(f^n(x), f^{n-m_n}(c_n)) \leq s_n \rightarrow 0$ where for big n we may assume that $c_n \in Q$. Hence $\omega(x) \subset \omega(Q) \cup (\cup_{q \in Q} \text{orb} q)$. Let us show that in fact this implies that $\omega(x) \subset \omega(Q)$. Indeed, otherwise there exist points $y \in \omega(x) \setminus \omega(Q)$ which then must belong to forward orbits of points of Q . Denote the set of all such points of Q by Q' ; clearly, $Q' = Q \setminus \omega(Q)$ and all points of $\omega(x) \setminus \omega(Q)$ are forward images of points of Q' . Now, every point $y \in \omega(x) \setminus \omega(Q)$ has at least one preimage in $\omega(x)$ (because $f|_{\omega(x)}$ is surjective) and this preimage cannot belong to $\omega(Q)$ (since $\omega(Q)$ is forward invariant and $y \notin \omega(Q)$ to begin with). Hence for every $q' \in Q'$ there exists $q'' \in \omega(x) \setminus \omega(Q)$ such that $f(q'') = q'$. The point q'' in turn is a forward image of some point of Q' under some iterate of f . Thus we can pull back the point q'' along this orbit and then repeat the argument. Since there are only finitely many points in the set Q' , at some moment the same point from Q' will have to appear twice which means that this is a periodic point. Yet this contradicts the fact that $Q' = Q \setminus \omega(Q)$ and shows that $\omega(x) = \omega(Q)$.

Let us show that if Q'' is the set of all recurrent points of Q then still $\omega(x) = \omega(Q'')$. Indeed, $Q \subset \omega(x) = \omega(Q)$. Hence if $q \in Q$ is not recurrent then there is a point $\hat{q} \neq q, \hat{q} \in Q$ such that $q \in \omega(\hat{q})$. Replace the set Q by the set $\hat{Q} = Q \setminus \{q\}$, then again

$\hat{Q} \subset \omega(x) = \omega(\hat{Q})$ and the argument can be repeated. After finitely many steps we get the set Q'' for which still $\omega(x) = \omega(Q'')$.

Now, let \tilde{Q} be the set of all points $\tilde{q} \in Q$ whose limit sets are maximal by inclusion among limit sets of points of Q . Let us show that all points of \tilde{Q} are recurrent. If $z \in \tilde{Q}$ is not recurrent then the fact that $Q \subset \omega(x) = \omega(Q)$ implies that there exists a point $z' \in Q$ such that $z \in \omega(z')$ and hence $\omega(z) \subsetneq \omega(z')$, a contradiction to the maximality of $\omega(z)$. So, $\tilde{Q} \subset Q''$ and clearly $\omega(x) = \omega(Q'') = \omega(\tilde{Q})$.

Let us show that there exists a point $t \in \tilde{Q}$ such that $\omega(x) = \omega(t)$. By way of contradiction assume that there is no such point in \tilde{Q} . To continue the argument we need to make a few choices of constants and points. Choose $\gamma > 0$ so that for any two points $y, z \in Q$ if $y \notin \omega(z)$ then $d(y, \omega(z)) > 2\gamma$ (this is possible since Q is finite). Then choose $\delta < \gamma$ so that for any two points u, v the fact that $d(u, v) < \delta$ implies $d(f(u), f(v)) < \gamma$ which is possible by continuity. Choose N so big that $s_n < \delta$ for $n > N$. Choose $n > N$ such that $c_n = c \in \tilde{Q}$. Since $\omega(c_n) \neq \omega(x)$ by the assumption then there exists the first moment $r > n$ such that $d(f^r(x), \omega(c)) > \delta$ (by the choice of γ there are points in $\omega(x)$ which are more than γ -distant from $\omega(c)$, e.g. those points of \tilde{Q} which do not belong to $\omega(c)$).

Consider the point c_r . Let us show that $c \in \omega(c_r)$ is impossible. Indeed, suppose that $c \in \omega(c_r)$. Then by the maximality of $\omega(c)$ we see that $\omega(c_r)$ is also maximal (so that $c_r \in \tilde{Q}$) and $\omega(c_r) = \omega(c)$. By the proven in the paragraph before the previous one, c_r is recurrent and since $d(f^r(x), f^{r-m_r}(c_r)) \leq s_r \leq \delta$ we see that $d(f^r(x), \omega(c)) \leq \delta$, a contradiction. So, $c \notin \omega(c_r)$. Vice versa, let us show that $c_r \notin \omega(c)$. Indeed, otherwise $d(f^r(x), f^{r-m_r}(c_r)) \leq s_n \leq \delta$ implies that again $d(f^r(x), \omega(c)) \leq \delta$, a contradiction.

Now, let us show that $m_r < r$. Indeed, otherwise $d(f^r(x), c_r) \leq s_r \leq \delta$. On the other hand $d(f^{r-1}(x), \omega(c)) \leq \delta$ by the choice of r . Thus $d(f^r(x), \omega(c)) \leq \gamma$ by the choice of δ . Together with $d(f^r(x), c_r) \leq \delta$ this implies that $d(c_r, \omega(c)) \leq \delta + \gamma < 2\gamma$ and hence $c_r \in \omega(c)$ by the choice of γ , a contradiction. So, $m_r < r$.

By the construction $d(f^{m_r}(x), c_r) \leq s_r \leq \delta$. Let us compare m_r with m_n and show by considering a few cases that all possibilities lead to a contradiction. First suppose that $m_r \geq m_n$. Then $d(f^{m_r}(x), \omega(c)) \leq \delta$ which together with $d(f^{m_r}(x), c_r) \leq \delta$ implies that $d(c_r, \omega(c)) \leq 2\delta < 2\gamma$ implying a above that $c_r \in \omega(c)$, a contradiction. Now, suppose that $m_r < m_n$. Then $d(f^{m_n-m_r}(c_r), f^{m_n}(x)) \leq s_r \leq \delta$ and $d(c, f^{m_n}(x)) \leq s_n \leq \delta$ imply that $d(f^{m_n-m_r}(c_r), c) < 2\gamma$ and hence by the choice of γ we have $c \in \omega(c_r)$, again the same contradiction. So, we show that the assumption that $\omega(c) \neq \omega(x)$ implies a contradiction which completes the proof of the fact that $\omega(c) = \omega(x)$. ■

To apply Theorems 10 and 11 we need Theorem 12 obtained in its final form in [BO] (see also [L] for polynomials with one critical point whose Julia sets are locally connected continua non-separating the plane, [BL2] for polynomials with any number of critical points whose Julia sets are locally connected continua non-separating the plane). To state Theorem 12 we need some information about closed invariant laminations (for the definitions see [Do], [McM], [Th]). Namely, if \sim is such laminations then we define an extension \simeq of \sim onto $\hat{\mathbb{C}} \setminus \mathbb{D}$ by declaring that a point in $\mathbb{C} \setminus \mathbb{D}$ is equivalent only to itself. Let $p : \mathbb{C} \setminus \mathbb{D} \rightarrow (\mathbb{C} \setminus \mathbb{D}) / \simeq$ be the factor map and denote $p(S^1)$ by J . Then J is a locally connected unshielded continuum, and since the map $\sigma : S^1 \rightarrow S^1$ defined by

$\sigma(z) = z^d$ acts on S^1 and the relation \sim is σ -invariant, we can consider a factor map $f : J \rightarrow J$. Notice, that the notions of a vertex and a critical point make sense for the map f and that there are examples of non-polynomial laminations (see [BL1]).

Theorem 12 [BO]. *Suppose that one of the assumptions below holds:*

- (1) \sim is an invariant lamination and $f : J \rightarrow J$ is the corresponding factor map;
- (2) f is a polynomial such that its Julia set J is locally connected.

Then $f|_J$ is backward stable.

Theorems 10, 11 and 12 imply Theorem 13 which in turn implies our Main Theorem.

Theorem 13. *Suppose that one of the assumptions below holds:*

- (1) \sim is an invariant lamination and $f : J \rightarrow J$ is the corresponding factor map;
- (2) f is a polynomial, J is its Julia set, and J is locally connected.

Then if x is a non-preperiodic non-precritical vertex of J then $\omega(x)$ coincides with $\omega(c(x))$ for a recurrent critical point $c(x)$ depending on x .

REFERENCES

- [BL1] A. Blokh, G. Levin, *An inequality for laminations, Julia sets and “growing trees”*, Erg. Th. and Dyn. Syst. (to appear).
- [BL2] ———, *Growing trees, laminations and the dynamics on the Julia set*, IHES Preprint IHES/M/99/77 (1999).
- [BM] A. Blokh, M. Misiurewicz, *Attractors for graph critical rational functions*, submitted to Trans. Amer. Math. Soc.
- [BO] A. Blokh, I. Oversteegen, *Backward stability for polynomial maps with locally connected Julia sets*, Preprint (2000).
- [BH] B. Branner, J. Hubbard, *The iteration of cubic polynomials. Part II: Patterns and parapatterns.*, Act Math. **169** (1992), 229-325.
- [CJY] L. Carleson, P. Jones, J.-C. Yoccoz, *Julia and John*, Bol. Soc. Bras. Mat. **25** (1994), 1–30.
- [CL] E. F. Collingwood, A. J. Lohwater, *The Theory of Cluster Sets*, Cambridge University Press, Cambridge, 1966.
- [Do] A. Douady, *Descriptions of compact sets in \mathbb{C}* , Topological methods in modern mathematics, Publish or Perish, 1993, pp. 429-465.
- [DH] A. Douady and J. H. Hubbard, *Etude dynamique des polynomes complexes I, II*, Publication Mathematiques d’Orsay **84-02** (1984); **85-04** (1985).
- [F] P. Fatou, *Sur les equations fonctionnelles*, Bul. Soc. Math. France **47** (1919), 161–271; **48** (1920), 33–94, 208–314.
- [H] J. H. Hubbard *Local connectivity of Julia sets and bifurcation loci: three theorems of J.-C. Yoccoz*, Topological methods in modern mathematics, Publish or Perish, 1993, pp. 467–511.
- [J] G. Julia, *Memoire sur l’iteration des fonctions rationnelles*, J. Math. Pure Appl. **8** (1919), 47–245.
- [Ki] J. Kiwi, *Rational rays and critical portraits of complex polynomials*, SUNY at Stony Brook and IMS Preprint #1997/15.
- [Kur] C. Kuratowski, *Topologie*, vol. 2, Warszawa-Wroclaw, 1950.
- [L] G. Levin, *On backward stability of holomorphic dynamical systems*, Fundamenta Mathematicae **158** (1998), 97–107.
- [McM] C. T. McMullen, *Complex dynamics and renormalization*, Annals of Mathematical Studies, vol. 135, Princeton University Press, Princeton, NJ, 1994.
- [Mi] J. Milnor, *Dynamics in One Complex Variable: Introductory Lectures*, SUNY at Stony Brook and IMS Preprint #1990/5.

- [Po] A. Poirier, *On post critically finite polynomials. Part two: Hubbard trees*, SUNY at Stony Brook and IMS Preprint #1993/7.
- [Sul] D. Sullivan, Conformal dynamical systems, in: *Geometric Dynamics*, Springer Lecture Notes, vol. 1007, Springer Verlag, NY, 1983, pp. 725–752.
- [Th] W. Thurston, *The combinatorics of iterated rational maps*, Preprint (1985).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA IN BIRMINGHAM, UNIVERSITY STATION,
BIRMINGHAM, AL 35294-2060
E-mail address: `ablokh@math.uab.edu`

INSTITUTE OF MATHEMATICS HEBREW UNIVERSITY GIVAT RAM 91904 JERUSALEM, ISRAEL
E-mail address: `levin@math.huji.ac.il`