
Bibliography

- [1] R. J. Adcock. Note on the method of least squares. *Analyst, London*, 4:183–184, 1877.
- [2] R. J. Adcock. A problem in least squares. *Analyst, London*, 5:53–54, 1878.
- [3] S. J. Ahn, W. Rauh, and H. J. Warnecke. Least-squares orthogonal distances fitting of circle, sphere, ellipse, hyperbola, and parabola. *Pattern Recog.*, 34:2283–2303, 2001.
- [4] A. Al-Sharadqah and N. Chernov. Error analysis for circle fitting algorithms. *Electr. J. Statist.*, 3:886–911, 2009.
- [5] Y. Amemiya and W. A. Fuller. Estimation for the nonlinear functional relationship. *Annals Statist.*, 16:147–160, 1988.
- [6] D. A. Anderson. The circular structural model. *J. R. Statist. Soc. B*, 27:131–141, 1981.
- [7] T. W. Anderson. Estimation of linear functional relationships: Approximate distributions and connections with simultaneous equations in econometrics. *J. R. Statist. Soc. B*, 38:1–36, 1976.
- [8] T. W. Anderson. Estimating linear statistical relationships. *Annals Statist.*, 12:1–45, 1984.
- [9] T. W. Anderson and H. Rubin. Statistical inference in factor analysis. In *Proc. Third Berkeley Symp. Math. Statist. Prob.*, volume 5, pages 111–150. Univ. California Press, 1956.
- [10] T. W. Anderson and T. Sawa. Exact and approximate distributions of the maximum likelihood estimator of a slope coefficient. *J. R. Statist. Soc. B*, 44:52–62, 1982.
- [11] I. O. Angell and J. Barber. An algorithm for fitting circles and ellipses to megalithic stone rings. *Science and Archaeology*, 20:11–16, 1977.
- [12] A. Atieg and G. A. Watson. Fitting circular arcs by orthogonal distance regression. *Appl. Numer. Anal. Comput. Math.*, 1:66–76, 2004.
- [13] J. Bentz. Private communication, 2001.
- [14] R. H. Berk. Sphericity and the normal law. *Annals Prob.*, pages 696–701, 1986.

- [15] M. Berman. Estimating the parameters of a circle when angular differences are known. *Appl. Statist.*, 32:1–6, 1983.
- [16] M. Berman. The asymptotic statistical behaviour of some estimators for a circular structural model. *Statist. Prob. Lett.*, 7:413–416, 1989.
- [17] M. Berman. Large sample bias in least squares estimators of a circular arc center and its radius. *CVGIP: Image Understanding*, 45:126–128, 1989.
- [18] M. Berman and D. Culpin. The statistical behavior of some least squares estimators of the center and radius of a circle. *J. R. Statist. Soc. B*, 48:183–196, 1986.
- [19] M. Berman and D. Griffiths. Incorporating angular information into models for stone circle data. *Appl. Statist.*, 34:237–245, 1985.
- [20] M. Berman and P. I. Somlo. Efficient procedures for fitting circles and ellipses with application to sliding termination measurements. *IEEE Trans. Instrum. Meas.*, 35:31–35, 1986.
- [21] R. H. Biggerstaff. Three variations in dental arch form estimated by a quadratic equation. *J. Dental Res.*, 51:1509, 1972.
- [22] P. T. Boggs, R. H. Byrd, J. E. Rogers, and R. B. Schnabel. Users reference guide for ODRPACK version 2.01. Technical Report NISTIR 4834, NIST, US Department of Commerce, 1992.
- [23] P. T. Boggs, R. H. Byrd, and R. B. Schnabel. A stable and efficient algorithm for nonlinear orthogonal distance regression. *SIAM J. Sci. Stat. Comput.*, 8:1052–1078, 1987.
- [24] F. L. Bookstein. Fitting conic sections to scattered data. *Computer Graphics and Image Processing*, 9:56–71, 1979.
- [25] J. A. Brandon and A. Cowley. A weighted least squares method for circle fitting to frequency response data. *J. Sound and Vibration*, 89:419–424, 1983.
- [26] D. Calvetti and L. Reichel. Gauss quadrature applied to trust region computations. *Numerical Algorithms*, 34:85–102, 2003.
- [27] R. J. Carroll, D. Ruppert, and L. A. Stefansky. *Measurement Error in Nonlinear Models*. Chapman & Hall, London, 1st edition, 1995.
- [28] R. J. Carroll, D. Ruppert, L. A. Stefansky, and C. M. Crainiceanu. *Measurement Error in Nonlinear Models: A Modern Perspective*. Chapman & Hall, London, 2nd edition, 2006.
- [29] G. Casella and R. L. Berger. *Statistical Inference*. Brooks/Cole, 3rd edition, 2002.
- [30] L. K. Chan and T. K. Mak. On the polynomial functional relationship. *J. R. Statist. Soc. B*, 47:510–518, 1985.

- [31] N. N. Chan. On circular functional relationships. *J. R. Statist. Soc. B*, 27:45–56, 1965.
- [32] Y. T. Chan, Y. Z. Elhalwagy, and S. M. Thomas. Estimation of circle parameters by centroiding. *J. Optimiz. Theory Applic.*, 114:363–371, 2002.
- [33] Y. T. Chan, B. H. Lee, and S. M. Thomas. Unbiased estimates of circle parameters. *J. Optimiz. Theory Applic.*, 106:49–60, 2000.
- [34] Y. T. Chan, B. H. Lee, and S. M. Thomas. Approximate maximum likelihood estimation of circle parameters. *J. Optimiz. Theory Applic.*, 125:723–733, 2005.
- [35] Y. T. Chan and S. M. Thomas. Cramer-Rao lower bounds for estimation of a circular arc center and its radius. *Graph. Models Image Proc.*, 57:527–532, 1995.
- [36] B. B. Chaudhuri and P. Kundu. Optimum circular fit to weighted data in multi-dimensional space. *Patt. Recog. Lett.*, 14:1–6, 1993.
- [37] C.-L. Cheng and A. Kukush. Non-existence of the first moment of the adjusted least squares estimator in multivariate errors-in-variables model. *Metrika*, 64:41–46, 2006.
- [38] C.-L. Cheng and J. W. Van Ness. On the unreplicated ultrastructural model. *Biometrika*, 78:442–445, 1991.
- [39] C.-L. Cheng and J. W. Van Ness. On estimating linear relationships when both variables are subject to errors. *J. R. Statist. Soc. B*, 56:167–183, 1994.
- [40] C.-L. Cheng and J. W. Van Ness. *Statistical Regression with Measurement Error*. Arnold, London, 1999.
- [41] N. Chernov. Fitting circles to scattered data: parameter estimates have no moments. *METRIKA*, *in press*, 2010.
- [42] N. Chernov and C. Lesort. Statistical efficiency of curve fitting algorithms. *Comp. Stat. Data Anal.*, 47:713–728, 2004.
- [43] N. Chernov and C. Lesort. Least squares fitting of circles. *J. Math. Imag. Vision*, 23:239–251, 2005.
- [44] N. Chernov and P. Sapirstein. Fitting circles to data with correlated noise. *Comput. Statist. Data Anal.*, 52:5328–5337, 2008.
- [45] N. I. Chernov and G. A. Ososkov. Effective algorithms for circle fitting. *Comp. Phys. Comm.*, 33:329–333, 1984.
- [46] W. Chojnacki, M. J. Brooks, and A. van den Hengel. Fitting surfaces to data with covariance information: fundamental methods applicable to computer vision. Technical Report TR99-03, Department of Computer Science, University of Adelaide, 1999. Available at

http://www.cs.adelaide.edu.au/~mjb/mjb_abstracts.html.

- [47] W. Chojnacki, M. J. Brooks, and A. van den Hengel. Rationalising the renormalisation method of Kanatani. *J. Math. Imaging & Vision*, 14:21–38, 2001.
- [48] W. Chojnacki, M. J. Brooks, A. van den Hengel, and D. Gawley. From FNS to HEIV: a link between two vision parameter estimation methods. *IEEE Trans. Pattern Analysis Machine Intelligence*, 26:264–268, 2004.
- [49] W. Chojnacki, M. J. Brooks, A. van den Hengel, and D. Gawley. FNS, CFNS and HEIV: A unifying approach. *J. Math. Imaging Vision*, 23:175–183, 2005.
- [50] W. R. Cook. On curve fitting by means of least squares. *Philos. Mag. Ser. 7*, 12:1025–1039, 1931.
- [51] I. D. Coope. Circle fitting by linear and nonlinear least squares. *J. Optim. Theory Appl.*, 76:381–388, 1993.
- [52] C. Corral and C. Lindquist. On implementing Kasa’s circle fit procedure. *IEEE Trans. Instrument. Measur.*, 47:789–795, 1998.
- [53] J. F. Crawford. A non-iterative method for fitting circular arcs to measured points. *Nucl. Instr. Meth.*, 211:223–225, 1983.
- [54] P. Delogne. Computer optimization of Deschamps’ method and error cancellation in reflectometry. In *Proc. IMEKO-Symp. Microwave Measurement (Budapest)*, pages 117–123, 1972.
- [55] W. E. Deming. The application of least squares. *Philos. Mag. Ser. 7*, 11:146–158, 1931.
- [56] G. R. Dolby. The ultrastructural relation: A synthesis of the functional and structural relations. *Biometrika*, 63:39–50, 1976.
- [57] I. Fazekas and A. Kukush. Asymptotic properties of an estimator in nonlinear functional errors-in-variables models with dependent error terms. *Computers Math. Appl.*, 34:23–39, 1997.
- [58] I. Fazekas and A. Kukush. Asymptotic properties of estimators in nonlinear functional errors-in-variables with dependent error terms. *J. Math. Sci.*, 92:3890–3895, 1998.
- [59] I. Fazekas, A. Kukush, and S. Zwanzig. Correction of nonlinear orthogonal regression estimator. *Ukr. Math. J.*, 56:1101–1118, 2004.
- [60] W. Feller. *An Introduction to Probability Theory and Its Applications*, Vol. 2. John Wiley & Sons, 2nd edition, 1971.
- [61] A. Fitzgibbon, M. Pilu, and R. Fisher. Direct least-square fitting of ellipses. In *International Conference on Pattern Recognition*, volume 21, pages 476–480, Vienna, August 1996.
- [62] A. W. Fitzgibbon and R. B. Fisher. A buyer’s guide to conic fitting. In

- British Machine Vision Conf.*, pages 513–522, 1995.
- [63] A. W. Fitzgibbon, M. Pilu, and R. B. Fisher. Direct least squares fitting of ellipses. *IEEE Trans. Pattern Analysis and Machine Intelligence*, 21:476–480, 1999.
- [64] J. L. Fleiss. The distribution of a general quadratic form in normal variables. *J. Amer. Statist. Assoc.*, 66:142–144, 1971.
- [65] P. R. Freeman. Note: Thom’s survey of the Avebury ring. *J. Hist. Astronom.*, 8:134–136, 1977.
- [66] W. A. Fuller. *Measurement Error Models*. L. Wiley & Son, New York, 1987.
- [67] W. Gander, G. H. Golub, and R. Strebler. Least squares fitting of circles and ellipses. *BIT*, 34:558–578, 1994.
- [68] J. Gates. Testing for circularity of spacially located objects. *J. Applied Statist.*, 20:95–103, 1993.
- [69] C.-F. Gauss. *Theoria Motus Corporum Coelestium in sectionibus conicis solem ambientium*. Book 2, 1809.
- [70] D. M. Gay. Computing optimal locally constrained steps. *SIAM J. Scientific. Statist. Computing*, 2:186–197, 1981.
- [71] P. E. Gill and W. Murray. Algorithms for the solution of the nonlinear least-squares problem. *SIAM J. Numer. Anal.*, 15:977–992, 1978.
- [72] C. Gini. Sull’interpolazione di una retta quando i valori della variabile indipendente sono affetti da errori accidentali. *Metron*, 1:63–82, 1921.
- [73] L. J. Gleser. Estimation in a multivariate ‘errors in variables’ regression model: Large sample results. *Annals Statist.*, 9:24–44, 1981.
- [74] L. J. Gleser. Functional, structural and ultrastructural errors-in-variables models. In *Proc. Bus. Econ. Statist. Sect. Am. Statist. Ass.*, pages 57–66, 1983.
- [75] L. J. Gleser. A note on G. R. Dolby’s unreplicated ultrastructural model. *Biometrika*, 72:117–124, 1985.
- [76] L. J. Gleser and G. S. Watson. Estimation of a linear transformation. *Biometrika*, 60:525–534, 1973.
- [77] G. H. Golub and C. F. Van Loan. An analysis of the total least squares problem. *SIAM J. Numer. Anal.*, 17:883–893, 1980.
- [78] G. H. Golub and C. F. van Loan. *Matrix computations*. J. Hopkins U. Press, 3rd edition, 1996.
- [79] Z. Griliches and V. Ringstad. Error-in-the-variables bias in nonlinear contexts. *Econometrica*, 38:368–370, 1970.
- [80] R. Halilović and J. Flusser. Numerically stable direct least squares fitting of

- ellipses. In *Sixth Conf. Cent. Europe Comput. Graph. Vis., WSCG'98, Conf. Proc.*, volume 1, pages 125–132, Plzen, Czech Rep., 1998.
- [81] R. Haliř and Ch. Menard. Diameter estimation for archaeological pottery using active vision. In Axel Pinz, editor, *Proc. 20th Workshop Austrian Assoc. Pattern Recognition (ÖAGM'96)*, pages 251–261, Schloss Seggau, Leibnitz, 1996.
- [82] M. Hansroul, H. Jeremie, and D. Savard. Fast circle fit with the conformal mapping method. *Nucl. Instr. Methods Phys. Res. A*, 270:498–501, 1988.
- [83] R. M. Haralick and L. G. Shapiro. *Computer and Robot Vision, Vol. I*. Addison-Wesley, Reading, MA, 1992.
- [84] <http://www.math.uab.edu/~chernov/cl>.
- [85] P. J. Huber. Robust estimation of a location parameter. *Annals Math. Statist.*, 35:73–101, 1964.
- [86] P. J. Huber. The behavior of maximum likelihood estimates under non-standard conditions. In *Proc. Fifth Berkeley Symp. Math. Stat. Prob.*, volume 1, pages 221–233, Univ. California Press, Berkeley, 1967.
- [87] P. J. Huber. *Robust Statistics*. Wiley-Interscience, 1981.
- [88] J. P. Imhof. Computing the distribution of quadratic forms in normal variables. *Biometrika*, 48:419–426, 1961.
- [89] W. N. Jessop. One line or two? *Appl. Statist.*, 1:131–137, 1952.
- [90] S. H. Joseph. Unbiased least-squares fitting of circular arcs. *Graph. Mod. Image Process.*, 56:424–432, 1994.
- [91] J. B. Kadane. Testing overidentifying restrictions when the disturbances are small. *J. Amer. Statist. Assoc.*, 65:182–185, 1970.
- [92] J. B. Kadane. Comparison of k-class estimators when the disturbances are small. *Econometrica*, 39:723–737, 1971.
- [93] K. Kanatani. Renormalization for unbiased estimation. In *Proc. 4th Intern. Conf. Computer Vision (ICCV'93)*, pages 599–606, Berlin, Germany, 1993.
- [94] K. Kanatani. Statistical bias of conic fitting and renormalization. *IEEE Trans. Pattern Analysis Machine Intelligence*, 16:320–326, 1994.
- [95] K. Kanatani. *Statistical Optimization for Geometric Computation: Theory and Practice*. Elsevier Science, Amsterdam, Netherlands, 1996.
- [96] K. Kanatani. Cramer-Rao lower bounds for curve fitting. *Graph. Mod. Image Process.*, 60:93–99, 1998.
- [97] K. Kanatani. For geometric inference from images, what kind of statistical method is necessary? *Mem. Faculty Engin. Okayama Univ.*, 37:19–28, 2002.

- [98] K. Kanatani. How are statistical methods for geometric inference justified? In *3rd Intern. Conf. Statist. Comput. Theories of Vision (SCTV 2003)*, 2003, Nice, France, 2003.
- [99] K. Kanatani. For geometric inference from images, what kind of statistical model is necessary? *Syst. Comp. Japan*, 35:1–9, 2004.
- [100] K. Kanatani. Optimality of maximum likelihood estimation for geometric fitting and the KCR lower bound. *Memoirs Fac. Engin. Okayama Univ.*, 39:63–70, 2005.
- [101] K. Kanatani. Unraveling the mystery of renormalization. *IPSJ SIG Technical Reports*, 38:15–22, 2005.
- [102] K. Kanatani. Ellipse fitting with hyperaccuracy. *IEICE Trans. Inform. Syst.*, E89-D:2653–2660, 2006.
- [103] K. Kanatani. Ellipse fitting with hyperaccuracy. In *Proc. 9th European Conf. Computer Vision (ECCV 2006)*, volume 1, pages 484–495, 2006.
- [104] K. Kanatani. Statistical optimization for geometric fitting: Theoretical accuracy bound and high order error analysis. *Int. J. Computer Vision*, 80:167–188, 2008.
- [105] Y. Kanazawa and K. Kanatani. Optimal line fitting and reliability evaluation. *IEICE Trans. Inform. Syst.*, E79-D:1317–1322, 1996.
- [106] V. Karimäki. Effective circle fitting for particle trajectories. *Nucl. Instr. Meth. Phys. Res. A*, 305:187–191, 1991.
- [107] V. Karimäki. Fast code to fit circular arcs. *Computer Physics Commun.*, 69:133–141, 1992.
- [108] I. Kåsa. A curve fitting procedure and its error analysis. *IEEE Trans. Inst. Meas.*, 25:8–14, 1976.
- [109] M. G. Kendall. Regression, structure, and functional relationships, Part I. *Biometrika*, 38:11–25, 1951.
- [110] M. G. Kendall. Regression, structure, and functional relationships, Part II. *Biometrika*, 39:96–108, 1952.
- [111] M. G. Kendall and A. Stuart. *The Advanced Theory of Statistics, Volume 2, Inference and Relationship*. Griffin, London, 4th edition, 1979.
- [112] D. Kincaid and W. Cheney. *Numerical Analysis*. Duxbury Press, 2nd edition, 2002.
- [113] T. C. Koopmans. *Linear regression analysis of economic time series*. De erven F. Bohn n.v., Haarlem, 1937.
- [114] A. Kukush, I. Markovsky, and S. Van Huffel. Consistent estimation in an implicit quadratic measurement error model. *Comput. Statist. Data Anal.*, 47:123–147, 2004.
- [115] A. Kukush, I. Markovsky, and S. Van Huffel. Consistent estimation of

- an ellipsoid with known center. In *COMPSTAT Proc. Comput. Statistics, 16th Symp., Prague, Czech Republic*, pages 1369–1376, Heidelberg, Germany, 2004. Physica-Verlag.
- [116] A. Kukush and E. O. Maschke. The efficiency of adjusted least squares in the linear functional relationship. *J. Multivar. Anal.*, 87:261–274, 2003.
- [117] C. H. Kummell. Reduction of observation equations which contain more than one observed quantity. *Analyst, London*, 6:97–105, 1879.
- [118] N. Kunitomo. Asymptotic expansions of the distributions of estimators in a linear functional relationship and simultaneous equations. *J. Amer. Statist. Assoc.*, 75:693–700, 1980.
- [119] U. M. Landau. Estimation of a circular arc center and its radius. *CVGIP: Image Understanding*, 38:317–326, 1987.
- [120] Y. Leedan and P. Meer. Heteroscedastic regression in computer vision: Problems with bilinear constraint. *Intern. J. Comp. Vision*, 37:127–150, 2000.
- [121] A.-M. Legendre. *Nouvelles méthodes pour la détermination des orbites des comètes*. Paris, 1806.
- [122] K. Levenberg. A method for the solution of certain non-linear problems in least squares. *Quart. Appl. Math.*, 2:164–168, 1944.
- [123] B. Lillekjendlie. Circular arcs fitted on a Riemann sphere. *Computer Vision Image Underst.*, 67:311–317, 1997.
- [124] K. B. Lim, K. Xin, and G. S. Hong. Detection and estimation of circular arc segments. *Pattern Recogn. Letters*, 16:627–636, 1995.
- [125] R. A. Liming. *Mathematics for Computer Graphics*. Aero Publishers, Fallbrook, CA, 1979.
- [126] D. V. Lindley. Regression lines and the linear functional relationship. *Suppl. J. R. Statist. Soc.*, 9:218–244, 1947.
- [127] A. Madansky. The fitting of straight lines when both variables are subject to error. *J. Amer. Statist. Ass.*, 54:173–205, 1959.
- [128] E. Malinvaud. *Statistical Methods of Econometrics*. North Holland Publ. Co, Amsterdam, 3rd edition, 1980.
- [129] K. V. Mardia and D. Holmes. A statistical analysis of megalithic data under elliptic pattern. *J. R. Statist. Soc. A*, 143:293–302, 1980.
- [130] I. Markovsky, A. Kukush, and S. Van Huffel. Consistent least squares fitting of ellipsoids. *Numerische Mathem.*, 98:177–194, 2004.
- [131] D. Marquardt. An algorithm for least squares estimation of nonlinear parameters. *SIAM J. Appl. Math.*, 11:431–441, 1963.
- [132] P. A. P. Moran. Estimating structural and functional relationships. *J.*

- Multivariate Anal.*, 1:232–255, 1971.
- [133] J. J. Moré. The Levenberg-Marquardt algorithm: Implementation and theory. In *Lecture Notes in Mathematics, Vol. 630*, pages 105–116, Berlin, 1978. Springer-Verlag.
- [134] J. J. Moré, B. B. Garbow, and K. E. Hillstom. User guide for MINPACK-1. Technical Report ANL-80-74, Argonne National Lab., IL (USA), 1980.
- [135] J. J. Moré and D. C. Sorensen. Computing a trust region step. *SIAM J. Scientif. Statist. Computing*, 4:553–572, 1983.
- [136] L. Moura and R. I. Kitney. A direct method for least-squares circle fitting. *Comp. Phys. Commun.*, 64:57–63, 1991.
- [137] Y. Nievergelt. Hyperspheres and hyperplanes fitted seamlessly by algebraic constrained total least-squares. *Linear Algebra Appl.*, 331:43–59, 2001.
- [138] Y. Nievergelt. A finite algorithm to fit geometrically all midrange lines, circles, planes, spheres, hyperplanes, and hyperspheres. *J. Numerische Math.*, 91:257–303, 2002.
- [139] N. Nussbaum. Asymptotic optimality of estimators of a linear functional relation if the ratio of the error variances is known. *Math. Op. Statist. Ser. Statist.*, 8:173–198, 1977.
- [140] M. O’Neill, L. G. Sinclair, and F. J. Smith. Polynomial curve fitting when abscissas and ordinates are both subject to error. *Computer J.*, 12:52–56, 1969.
- [141] G. A. Ososkov. P10-83-187 (in Russian). Technical report, JINR Dubna, 1983.
- [142] W. M. Patefield. On the validity of approximate distributions arising in fitting linear functional relationship. *J. Statist. Comput. Simul.*, 5:43–60, 1976.
- [143] W. M. Patefield. Multivariate linear relationships: Maximum likelihood estimation and regression bounds. *J. R. Statist. Soc. B*, 43:342–352, 1981.
- [144] K. Pearson. On lines and planes of closest fit to systems of points in space. *Phil. Mag.*, 2:559–572, 1901.
- [145] S.-C. Pei and J.-H. Horng. Optimum approximation of digital planar curves using circular arcs. *Pattern Recogn.*, 29:383–388, 1996.
- [146] W. A. Perkins. A model-based vision system for industrial parts. *IEEE Trans. Comput.*, 27:126–143, 1978.
- [147] M. Pilu, A. Fitzgibbon, and R. Fisher. Ellipse-specific direct least-square fitting. In *IEEE International Conference Image Processing*, Lausanne,

September 1996.

- [148] M. J. D. Powell. A hybrid method for nonlinear equations. In P. Robinson, editor, *Numerical Methods for Nonlinear Programming Algebraic Equations*, pages 87–144, London, 1970. Gordon and Beach Science.
- [149] M. J. D. Powell. A new algorithm for unconstrained optimization. In J. B. Rosen, O. L. Mangasarian, and K. Ritter, editors, *Nonlinear Programming, Proc. Sympos. Math. Res. Center*, pages 31–66, New York, 1970.
- [150] V. Pratt. Direct least-squares fitting of algebraic surfaces. *Computer Graphics*, 21:145–152, 1987.
- [151] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery. *Numerical Recipes in C++*. Cambridge Univ. Press., 2nd edition, 2002.
- [152] P. Rangarajan and K. Kanatani. Improved algebraic methods for circle fitting. *Electr. J. Statist.*, 3:1075–1082, 2009.
- [153] H. Robbins and J. Pitman. Application of the method of mixtures to quadratic forms in normal variates. *Annals Math. Statist.*, 20:552–560, 1949.
- [154] J. Robinson. The distribution of a general quadratic form in normal variables. *Austral. J. Statist.*, 7:110–114, 1965.
- [155] S. M. Robinson. Fitting spheres by the method of least squares. *Commun. Assoc. Comput. Mach.*, 4:491, 1961.
- [156] C. F. Roos. A general invariant criterion of fit for lines and planes where all variates are subject to error. *Metron*, 13:3–30, 1937.
- [157] C. Rorres and D. G. Romano. Finding the center of a circular starting line in an ancient Greek stadium. *SIAM Rev.*, 39:745–754, 1997.
- [158] P. L. Rosin and G. A. W. West. Segmentation of edges into lines and arcs. *Image Vision Comp.*, 7:109–114, 1989.
- [159] C. Rusu, M. Tico, P. Kuosmanen, and E. J. Delp. Classical geometrical approach to circle fitting – review and new developments. *J. Electron. Imaging*, 12:179–193, 2003.
- [160] Ed. S. van Huffel. *Recent advances in total least squares techniques and errors-in-variables modeling*. SIAM, 1997.
- [161] Ed. S. van Huffel. *Total Least Squares and Errors-in-Variables Modeling*. Kluwer, Dordrecht, 2002.
- [162] R. Safaee-Rad, I. Tchoukanov, B. Benhabib, and K. C. Smith. Accurate parameter estimation of quadratic curves from grey-level images. *CVGIP: Image Understanding*, 54:259–274, 1991.
- [163] P. D. Sampson. Fitting conic sections to very scattered data: an iterative

- refinement of the Bookstein algorithm. *Comp. Graphics Image Proc.*, 18:97–108, 1982.
- [164] B. Sarkar, L. K. Singh, and D. Sarkar. Approximation of digital curves with line segments and circular arcs using genetic algorithms. *Pattern Recogn. Letters*, 24:2585–2595, 2003.
- [165] E. Saund. Identifying salient circular arcs on curves. *CVGIP: Image Understanding*, 58:327–337, 1993.
- [166] C. Shakarji. Least-squares fitting algorithms of the NIST algorithm testing system. *J. Res. Nat. Inst. Stand. Techn.*, 103:633–641, 1998.
- [167] S. Shklyar, A. Kukush, I. Markovsky, and S. Van Huffel. On the conic section fitting problem. *J. Multivar. Anal.*, 98:588–624, 2007.
- [168] M. E. Solari. ‘Maximum Likelihood solution’ of the problem of estimating a linear functional relationship. *J. R. Statist. Soc. B*, 31:372–375, 1969.
- [169] P. I. Somlo and J. D. Hunter. A six-port reflectometer and its complete characterization by convenient calibration procedures. *IEEE Trans. Microwave Theory Tech.*, 30:186–192, 1982.
- [170] H. Späth. Least-squares fitting by circles. *Computing*, 57:179–185, 1996.
- [171] H. Späth. Orthogonal least squares fitting by conic sections. In *Recent Advances in Total Least Squares techniques and Errors-in-Variables Modeling*, pages 259–264. SIAM, 1997.
- [172] T. Steihaug. Conjugate gradient method and trust regions in large scale optimization. *SIAM J. Numer. Anal.*, 20:626–637, 1983.
- [173] A. Strandlie and R. Frühwirth. Error analysis of the track fit on the Riemann sphere. *Nucl. Instr. Methods Phys. Res. A*, 480:734–740, 2002.
- [174] A. Strandlie, J. Wroldsen, and R. Frühwirth. Treatment of multiple scattering with generalized Riemann sphere track fit. *Nucl. Instr. Methods Phys. Res. A*, 488:332–341, 2002.
- [175] A. Strandlie, J. Wroldsen, R. Frühwirth, and B. Lillekjendlie. Particle tracks fitted on the Riemann sphere. *Computer Physics Commun.*, 131:95–108, 2000.
- [176] G. Taubin. Estimation of planar curves, surfaces and nonplanar space curves defined by implicit equations, with applications to edge and range image segmentation. *IEEE Trans. Pattern Analysis Machine Intelligence*, 13:1115–1138, 1991.
- [177] A. Thom. A statistical examination of the megalithic sites in Britain. *J. Royal Statist. Soc. A*, 118:275–295, 1955.
- [178] A. Thom and A. S. Thom. A megalithic lunar observatory in Orkney:

- the ring of Brogar and its cairns. *J. Hist. Astronom.*, 4:111–123, 1973.
- [179] A. Thom, A. S. Thom, and T. R. Foord. Avebury (1): a new assessment of the geometry and metrology of the ring. *J. Hist. Astronom.*, 7:183–192, 1976.
- [180] S. M. Thomas and Y. T. Chan. A simple approach for the estimation of circular arc center and its radius. *CVGIP: Image Understanding*, 45:362–370, 1989.
- [181] B. Triggs. A new approach to geometric fitting, ICCV'98, Bobbay (unpublished), 1998.
- [182] B. Triggs. Optimal estimation of matching constraints. In *3D Structure from Multiple Images of Large-Scale Environments*, volume 1506, pages 63–77, Berlin/Heidelberg, 1998. Springer.
- [183] K. Turner. *Computer perception of curved objects using a television camera*. PhD thesis, University of Edinburgh, 1974. Dept. of Machine Intelligence.
- [184] D. Umbach and K. N. Jones. A few methods for fitting circles to data. *IEEE Trans. Instrument. Measur.*, 52:1181–1885, 2003.
- [185] S. van Huffel and J. Vandewalle. *The Total Least Squares Problem: Computational Aspects and Analysis*. SIAM, Philadelphia, 1991.
- [186] G. Vosselman and R. M. Haralick. Performance analysis of line and circle fitting in digital images. In *Workshop on Performance Characteristics of Vision Algorithms*, April 1996.
- [187] A. Wald. The fitting of straight lines if both variables are subject to error. *Ann. Math. Statist.*, 11:284–300, 1940.
- [188] C. M. Wang and C. T. Lam. A mixed-effects model for the analysis of circular measurements. *Technometrics*, 39:119–126, 1997.
- [189] M. Werman and D. Keren. A Bayesian method for fitting parametric and nonparametric models to noisy data. *IEEE Trans. Pattern Analysis and Machine Intelligence*, 23:528–534, 2001.
- [190] M. E. Whalen. Ceramic vessel size estimation from sherds: An experiment and a case study. *J. Field Archaeology*, 25:219–227, 1998.
- [191] J. Wolfowitz. Estimation of structural parameters when the number of incidental parameters is unbounded. *Annals Math. Statist.*, 25:811, 1954.
- [192] K. M. Wolter and W. A. Fuller. Estimation of nonlinear errors-in-variables models. *Annals Statist.*, 10:539–548, 1982.
- [193] Z. Wu, L. Wu, and A. Wu. The robust algorithms for finding the center of an arc. *Comp. Vision Image Under.*, 62:269–278, 1995.
- [194] S. J. Yin and S. G. Wang. Estimating the parameters of a circle by

- heteroscedastic regression models. *J. Statist. Planning Infer.*, 124:439–451, 2004.
- [195] P. C. Yuen and G. C. Feng. A novel method for parameter estimation of digital arcs. *Pattern Recogn. Letters*, 17:929–938, 1996.
- [196] E. Zelniker and V. Clarkson. A statistical analysis of the Delogne-Kåsa method for fitting circles. *Digital Signal Proc.*, 16:498–522, 2006.
- [197] S. Zhang, L. Xie, and M. D. Adams. Feature extraction for outdoor mobile robot navigation based on a modified gaussnewton optimization approach. *Robotics Autonom. Syst.*, 54:277–287, 2006.
- [198] Z. Zhang. Parameter estimation techniques: A tutorial with application to conic fitting. *Inter. J. Image & Vision Comp.*, 15:59–76, 1997.