Name:

Signature:

Exam IV Calculus I; Fall 2009

Part I

Part I consists of 10 questions, each worth 5 points. Clearly show your work for each of the problems listed.

(1) Let $f(x) = 4x^3 - 12x + 2$. Find all local max/min of f(x). State both x and y coordinates. Answer: local max at (-1, 10) and local min at (1, -6).

(2) Find the absolute max/min of $f(x) = 10 - x^2$ on the interval [-1, 2]. Give both x and y-coordinates and justify your answer. Answer: absolute max at (0, 10) and absolute min at (2, -6).

(3) Find two positive numbers whose product is 100 and whose sum is minimal. (You must justify your answer.) Answer: 10 and 10. (4) Let $f'(x) = (x-2)^2(x-1)(x+1)$. Note that you are already given the derivative f'(x). Find all critical points, where f(x) is increasing and decreasing, and also find the xcoordinate(s) of all local max/min. Answer: increasing on $(-\infty, -1)$ and $(1, \infty)$, decreasing on (-1, 1). Local min at x = 1, local max at x = -1.

(5) If $f''(x) = (x-1)^2(x+3)$ find where f(x) is concave up and where it is concave down. Also find all points of inflection. Note that you are given f''(x)! Answer: concave up on $(-3, \infty)$, concave down on $(-\infty, -3)$, inflection point at x = -3.

(6) Find the most general **anti-**derivative of $\frac{3-x+4\sqrt{x}}{x^3}$. Answer: $-\frac{3}{2}x^{-2} + x^{-1} - \frac{8}{3}x^{-3/2} + C$. (7) Find the most general **anti-**derivative of $\cos(x) - \frac{1}{x}$. Answer: $\sin x - \ln |x| + C$.

(8) Find all asymptotes of the function $\frac{x^{3}+5}{x(x-2)(x+1)}$. Answer: vertical: x = 0, x = 2, x = -1 and horizontal y = 1.

(9) If the acceleration is given by a(t) = 6t, v(0) = 2 and s(0) = 1, find an expression for the position s(t). Answer: $s(t) = t^3 + 2t + 1$.

(10) If $f(x) = x^3$ find the number c that satisfies the conclusion of the mean value theorem on the interval [0, 2]. Answer: $c = 2/\sqrt{3}$.

Part II

Part II consists of 3 problems; the number of points for each part are indicated by [x pts]. You must show the relevant steps (as we did in class) and justify your answer to earn credit. Simplify your answer when possible.

(1) **[15 pts]** Find the absolute max/min of the function $f(x) = (x^2 - 1)^3$ on the interval [-2, 3]. Answer: absolute min (0, -1), absolute max (3, 512).

(2) Given the function $f(x) = \frac{x^2-9}{x^2-1}$,

(a) [2 pts] Find the x and y intercepts of the function. Answer: $x = \pm 3$ and y = 9.

(b) [3 pts] Find all asymptotes. Answer: vertical x = 1 and x = -1, horizontal y = 1. (c) [4 pts] Find the open intervals where f(x) is increasing and the open intervals where f(x) is decreasing, Answer: increasing on (0,1) and $(1,\infty)$, decreasing on $(-\infty, -1)$ and (-1, 0).

(d) [2 pts] Find the local maximum and local minimum value(s) of f(x). (Be sure to give the x and y coordinate of each of them).

Answer: local min at (0, 9), no local max.

(e) [2 pts] Find all open intervals where the graph of f(x) is concave up and all open intervals where the graph is concave down.
Answer: concave up on (-1, 1), concave down on (-∞, -1) and (1, ∞).

(f) [2 pts] Find all points of inflection (be sure to give the x and y coordinate of each point). Answer: x = 1 and x = -1.

(g) [6 pts] Use the above information to graph the function on the next page. Indicate all relevant information in the graph. Put the graph of Problem 2 on this page.

(3) **[14 pts]** A drilling rig in the ocean is 10 mi of shore. A refinery is located along the coast 15 mi away from the point on the shore closest to the rig. If under water pipe lines cost \$5 per mi and land-based pipe costs \$4 per mi, what is the least expensive way to run the line.



Partial answer: The total cost of the pipe is $f(x) = 4(15-x) + 5\sqrt{10^2 + x^2}$ you need to find x where this function is a minimum.