

1. Use the Gershgorin theorem to show that a symmetric, strictly row diagonally dominant real matrix with positive diagonal elements is positive definite.
2. (JPE, May 1989) Let A be a real symmetric matrix, with eigenvalues λ_i , $1 \leq i \leq n$, satisfying

$$|\lambda_1| > |\lambda_2| \geq \cdots \geq |\lambda_n|$$

If x_1 is an eigenvector corresponding to λ_1 , and z_0 is a vector satisfying $z_0^T x_1 \neq 0$, prove that

$$\lim_{k \rightarrow \infty} \frac{z_0^T A^k z_0}{z_0^T A^{k-1} z_0} = \lambda_1$$

3. (JPE, May 2003) Let A be a symmetric matrix with eigenvalues such that $|\lambda_1| > |\lambda_2| \geq \cdots \geq |\lambda_{n-1}| > |\lambda_n|$. Suppose $z \in \mathbb{R}^n$ with $z^T x_1 \neq 0$, where $Ax_1 = \lambda_1 x_1$. Prove that, for some constant C ,

$$\lim_{k \rightarrow \infty} \frac{A^k z}{\lambda_1^k} = Cx_1$$

and use this result to devise a reliable algorithm for computing λ_1 and x_1 . Explain how the calculation should be modified to obtain (a) λ_n and (b) the eigenvalue closest to 2.

4. (JPE, September 1996) The matrix

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 2 & 1 \end{pmatrix}$$

has eigenpairs

$$(\lambda, x) = \left(2, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right), \left(-1, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right), \left(3, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right),$$

Suppose the power method is applied with starting vector

$$z_0 = [1, 1, -1]^T / \sqrt{3}$$

- (a) Determine whether or not the iteration will converge to an eigenpair of A , and if so, which one. Assume exact arithmetic.
- (b) Repeat (a), except now use the inverse iteration with the same starting vector z_0 and the Rayleigh quotient of z_0 as approximation for the eigenvalue.
- (c) Now answer both (a) and (b) again, except this time use standard fixed precision floating point arithmetic, i.e. computer arithmetic.